

ANNEX F
INVASIVE AND NUISANCE SPECIES MANAGEMENT PLAN

This page intentionally left blank.

Table of Contents

F	INVASIVE AND NUISANCE SPECIES MANAGEMENT PLAN	F-1
F.1	Introduction.....	F-1
F.2	Status of Priority Species and Their Impacts.....	F-3
F.2.1	Plants	F-3
F.2.1.1	Widely Established Species.....	F-3
F.2.1.2	Australian pine.....	F-3
F.2.1.3	Brazilian pepper	F-4
F.2.1.4	Cogongrass	F-4
F.2.1.5	Dwarf rotala.....	F-4
F.2.1.6	Hydrilla	F-5
F.2.1.7	Melaleuca	F-5
F.2.1.8	Old World Climbing Fern	F-5
F.2.1.9	Shoebuttan ardesia	F-5
F.2.1.10	Torpedograss	F-6
F.2.1.11	Water Hyacinth.....	F-6
F.2.1.12	Water Lettuce.....	F-6
F.2.1.13	Water primrose	F-6
F.2.1.14	Cattail	F-7
F.2.1.15	Localized/Potential EDRR Species.....	F-7
F.2.1.15.1	Tropical American Watergrass.....	F-7
F.2.1.15.2	Mile-a-minute	F-7
F.2.1.15.3	Tropical Nutrush	F-8
F.2.1.15.4	Wright’s Nutrush	F-8
F.2.2	Animals	F-8
F.2.2.1	Widely Established Species.....	F-8
F.2.2.1.1	Redbay Ambrosia Beetle (laurel wilt).....	F-8
F.2.2.1.2	Asian Swamp Eel.....	F-9
F.2.2.1.3	Cuban Treefrog.....	F-9
F.2.2.1.4	Feral Hog.....	F-10
F.2.2.1.5	Green Iguana	F-11
F.2.2.1.6	Purple Swamphen.....	F-11
F.2.2.1.7	Island Applesnail.....	F-12
F.2.2.1.8	Burmese Python	F-12
F.2.2.1.10	Veiled Chameleon	F-12
F.2.2.2	Localized/Early Detection Rapid Response Species	F-13
F.2.2.2.1	Northern African Python.....	F-13
F.2.2.2.2	Argentine Black and White Tegu.....	F-13
F.2.2.2.3	Nile Monitor	F-14
F.2.2.2.4	Spectacled Caiman.....	F-14
F.2.2.2.5	Giant African Land Snail.....	F-14
F.3	Introduction to Invasive Species Management	F-15

F.3.1	Prevention	F-15
F.3.2	Education and Public Awareness.....	F-15
F.3.3	Monitoring.....	F-15
F.3.4	Early Detection and Rapid Response.....	F-16
F.3.5	Control and Management.....	F-16
F.3.6	Risk and Uncertainties Related to Invasive Species	F-17
F.4	Existing Management Programs	F-18
F.4.1	South Florida Water Management District.....	F-18
F.4.2	US Army Corps of Engineers.....	F-19
F.4.3	U.S. Department of Agriculture / University of Florida.....	F-19
F.4.4	Florida Fish and Wildlife Conservation Commission	F-19
F.4.5	Invasive Animals	F-19
F.5	Existing Monitoring Programs.....	F-20
F.6	Management Strategy and Plan.....	F-20
F.6.1	Surveillance – Early Detection and Rapid Response	F-21
F.6.2	Control.....	F-22
F.6.3	Monitoring.....	F-23
F.6.4	Pre-construction Phase	F-23
F.6.5	Design and Construction Phases	F-23
F.6.6	Operational Testing and Monitoring Period	F-25
F.6.7	OMRR&R Phase	F-25
F.6.8	Specific Control by Project Feature – Construction Phase	F-26
F.6.8.1	Region 1.....	F-26
North Feeder STA.....		F-26
F.6.8.2	Region 2.....	F-27
Wingate Mill Canal Plug		F-27
Wingate Mill & Lard Can Canal Modifications/Discharge Slough		F-27
F.6.8.3	Region 3.....	F-28
L-28i Inline Weir.....		F-28
L-28i Levee Degrade and Canal Backfill.....		F-28
Restoration Areas		F-29
F.6.8.4	Region 4.....	F-29
L-28(S) Levee Degrade, Canal Backfill and Gated Culverts		F-29
L-29 Plug		F-30
Culverts – 11 Mile Road, US 41 (Tamiami Trail) and Loop Road		F-30
F.6.9	Specific Control by Project Feature – OMRR&R Phase.....	F-31
F.6.9.1	Region 1.....	F-31
North Feeder STA.....		F-31
F.6.9.2	Region 2.....	F-31
Wingate Mill Canal Plug		F-31

Wingate Mill & Lard Can Canal Modifications/Discharge Slough F-32
 F.6.9.3 Region 3..... F-32
 L-28i Inline Weir..... F-32
 L-28i Levee Degrade and Canal Backfill..... F-32
 Restoration Areas F-32
 F.6.9.4 Region 4..... F-33
 L-29 Plug F-33
F.7 Education / Outreach Opportunities at Recreational Areas.....F-33
F.8 CostsF-34
 - **F-39**
F.9 REFERENCES.....F-50

List of Tables

Table F-1. Invasive Plant Species Documented in the Project Area F-35
 Table F-2. Invasive Plant Species Documented in the Project Area (Totals from Table F-1
 above). F-38
 Table F-3. Invasive Animal Species Documented in the Project Area. F-39
 Table F-4. Invasive Animal Species Documented in the Project Area (Totals from Table F-3
 Above) F-41
 Table F-5. WERP Invasive and Nuisance Species Management and Monitoring Costs. F-42
 Table F-6. Invasive and Nuisance Species Management Costs – Construction PHASE F-43
 Table F-7. Invasive and Nuisance Species Management Costs – OMRR&R Phase..... F-46
 Table F-8. Invasive and Nuisance Species Management Costs F-49

This page intentionally left blank.

F INVASIVE AND NUISANCE SPECIES MANAGEMENT PLAN

In accordance with the Comprehensive Everglades Restoration Plan (CERP) Guidance Memorandum 062.00 (CGM62), Invasive Species, the WERP will incorporate invasive and nuisance species assessments and management of those species into pertinent planning documents and phases of the project. The Invasive and Nuisance Species Management Plan (INSMP) is a living document and will be updated throughout the Design, Construction and Operations, Maintenance, Repair, Replacement, and Rehabilitation (OMRR&R) phases.

The Project Partnership Agreement (PPA) and the Construction Phasing, Transfer, and Warranty (CPTW) Plan are developed and agreed to prior to construction. The documents outline the responsibilities of the federal and non-federal sponsor during the construction phase, the operational testing and monitoring period, and the OMRR&R phase, and will include the cost estimates associated with this INSMP. This INSMP shall be included with the CPTW Plan.

F.1 Introduction

WERP will reestablish sheet flow from the northern portion of the study area, across the Seminole Tribe of Florida Big Cypress Reservation, into the Big Cypress National Preserve (BCNP) and along historic flow paths toward the southern coast of Florida. The WERP study incorporates three components of the CERP which include Big Cypress / L 28 Interceptor Modifications (Component CCC), WCA 3 Decompartmentalization and Sheetflow Enhancement (Component QQ), and flow to the northwest and central Water Conservation 3A (Component RR).

The WERP study area includes ~772,700 acres and focuses on the “western flow way” of the Everglades. It is approximately ~1,200 square miles due to the broad, relatively flat topography of the watershed and the basins that provide inflows to the L-28 canals. The L-1 Canal is the northern boundary of the project area. The L-2 Canal, Stormwater Treatment Area (STA) 5/6, and the eastern boundary of the Miccosukee Tribe of Indians of Florida Alligator Alley Reservation delineate the eastern boundary. Hydrologic basin boundaries mark the western edge of the study area. The southern boundary encompasses portions of US Highway 41, Loop Road, and a southern Miccosukee Tribe of Indians of Florida (Alligator Alley) Reservation area.

Nationally, more than 50,000 species of introduced plants, animals, and microbes cause more than \$120 billion in economic damages and control costs each year (Pimentel et al. 2005). Not all introduced species become invasive species. According to the US Congress Office of Technology Assessment “Harmful Non-indigenous Species in the United States” report, approximately 10 to 15% of introduced species will become established and 10% of the established species may become invasive.

Executive Order (E.O.) 13112, entitled Invasive Species, signed February 3, 1999, states an “invasive species means an alien species whose introduction does or is likely to cause economic or environmental harm or harm to human health.” Alien species means, with respect to a particular ecosystem, any species, including its seeds, eggs, spores, or other biological material capable of propagating that species, that is not native to that ecosystem. Invasive species are broadly defined and can be a plant, animal, fungus, plant disease, livestock disease or other organism. The terms ‘alien’ and ‘exotic’ also refer to non-native species. A native species is defined as a species that historically occurred or currently occurs in a particular ecosystem and is not the result of an introduction.

Invasive non-native species decrease biodiversity, displace native plant and animal communities, reduce wildlife habitat and forage opportunities, alter the rates of soil erosion and accretion, alter fire regimes, upset predator/prey relationships, alter hydrology, degrade environmental quality and spread diseases to native plants, animals and other organisms. Furthermore, invasive species are the second largest threat to biodiversity following only habitat destruction (Wilcove et al. 1998); invasive species are second in destructive nature only to human development. In the United States, invasive species directly contributed to the decline of 49% of the T&E species (Wilcove et al. 1998). In addition to environmental impacts, invasive species impact human health, reduce agricultural production and property values, degrade aesthetic quality, decrease recreational opportunities and threaten the integrity of human infrastructure such as waterways/navigation channels, locks, levees, dams and water control structures.

Florida is particularly vulnerable to the introduction, invasion and naturalization of non-native species. This is due to several factors including a subtropical climate, dense human population centers, major ports of entry and the pet, aquarium and ornamental plant industries. Major disturbance to the landscape has also increased Florida's vulnerability to invasive species. Alteration of the landscape for urban development, flood control and agricultural uses has exacerbated non-native plant and animal invasions. Florida is listed as one of the states with the largest number of invasive species. This list also includes Hawaii, California, and Louisiana. On average, 10 new organisms per year are introduced into Florida that are capable of establishing and becoming invasive and causing environmental harm. Approximately 90% of the plants and animals that enter the continental United States enter through the port of Miami (Cuda 2009a). Stein, Kutner and Adams (2000) estimated that over 32,000 exotic species (25,000 plants and 7,000 animals) have been introduced into Florida. There are approximately 4,000-5,000 native species of plants and animals in Florida. The number of non-native species that have been introduced is eight times the total number of native species in the entire state.

Significant scientific evidence and research document invasive non-native plants are degrading and damaging south Florida natural ecosystems (Doren and Ferriter 2001). Many species are causing significant ecological impacts by crowding out and displacing native plants, altering soil types and soil/water chemistry, altering ecosystem functions such as carbon sequestration, nutrient cycling and fire regimes and reducing gene pools and genetic diversity. Non-native invasive animal distribution, extent and impacts are not well understood, however implications of invasive animals are apparent in south Florida. It has been documented there are 14 non-native species that are causing direct impacts to threatened and endangered species and rare habitats. Holm et al. (1977) documented that 19 species within Florida are among the world's worst weeds. It is estimated that federal, state, and county agencies in Florida spend between \$94 million and \$127 million each year in an effort to manage invasive non-native plants (GAO 2000).

Invasive species are a major threat to the success of CERP. "The intent of CERP is to restore, preserve, and protect the south Florida ecosystem while providing for other water-related needs of the region. CERP focuses on hydrologic restoration to improve degraded natural habitat in the south Florida ecosystem. Hydrologic restoration alone cannot ensure habitat restoration" (USACE and SFWMD 2010). In order to restore the Everglades and ensure south Florida's natural ecosystems are preserved and remain intact, invasive species must be comprehensively addressed (Doren and Ferriter 2001). The lack of management will allow invasive non-native species to flourish and to continue to out-compete native species.

F.2 Status of Priority Species and Their Impacts

Information regarding both plants and animals is presented below in three categories: widely established within the project area, localized/potential early detection rapid response (EDRR) species and other species of concern.

F.2.1 Plants

According to the Atlas of Florida Plants, 4,875 plant species have been documented Florida. Of the 4,875 plant species, 1,557 were considered non-native and were naturalized (freely reproducing) populations. The Florida Exotic Pest Plant Council (FLEPPC) identifies 79 of the 1,557 species of non-native plants as Category I and 86 as Category II species in the 2021 Invasive Plant List. Searches through existing data and resources indicate 79 non-native plant species have been documented to occur within the project area (**Table F-1** and **Table F-2**). Other non-native species are probably present; however, documented citations could not be located. Of the 79 species of plants documented to occur within the project area, there are 39 FLEPPC Category I species, 19 FLEPPC Category II species, and 13 Florida Noxious Weed species.

A primary native nuisance species within the project area is cattail (*Typha* spp.). Many areas within the project area have been invaded by cattails. This is attributed to water with increased phosphorus being delivered to these areas which began in the late 1950s. Areas where water control structures, conveyance features and levees exist provide a suitable habitat for invasion and expansion of cattail.

F.2.1.1 Widely Established Species

Plants that are widely established within the project area that are managed for long term suppression include Australian pine (*Casuarina equisetifolia*), Brazilian pepper (*Schinus terebinthifolius*), cogongrass (*Imperata cylindrical*), creeping water-primrose (*Ludwigia* spp.), dwarf rotalla (*Rotala rotundifolia*), hydrilla (*Hydrilla verticillata*), melaleuca (*Melaleuca quinquenervia*), Old World climbing fern (*Lygodium microphyllum*), torpedo grass (*Panicum repens*), shoe button ardisia (*Ardisia crenata*), water hyacinth (*Eichhornia crassipes*), water lettuce (*Pistia stratiotes*), and cattail (*Typha* spp.).

Summaries on the distribution and impacts of these widely established species are included below. Other non-native plant species with limited or localized distributions or which have a high potential to spread into the project area are also discussed.

F.2.1.2 Australian pine

Australian pine is an evergreen tree that can grow to 150 feet tall. It has inconspicuous flowers and produces tiny fruit, a 1-seeded, winged nutlet that is formed in a woody cone-like cluster. Australian pine is a prolific seed producer and seeds are dispersed by birds, wind and water flow. It is native to Australia, the south Pacific Islands, and Southeast Asia. Australian pine was introduced in the late 1800s and was planted extensively in south Florida as windbreaks and shade trees. It inhabits sandy shores and pinelands and is salt tolerant. It also invades disturbed sites such as filled wetlands, roadsides, cleared undeveloped land, canal banks, and levees. Australian pine grows rapidly, shading out native species. It produces dense litter accumulation, causes beach erosion, and produces an allelopathic agent that inhibits growth of other species. It also interferes with nesting of sea turtles and the American crocodile (Langeland and Burks 1998).

F.2.1.3 Brazilian pepper

Brazilian pepper is an evergreen shrub or tree that can grow up to 40 feet tall. It forms dense thickets and is a prolific seed producer. It produces a small bright red fruit in the form of a spherical drupe. Brazilian pepper is native to Brazil, Argentina, and Paraguay and was imported in the 1840s as an ornamental plant (Langeland and Burks 1998). Brazilian pepper inhabits natural areas such as pinelands, hardwood hammocks and mangrove forests. It is an aggressive pioneer species that quickly colonizes and thrives in disturbed areas (Francis n.d.) such as fallow farmland, fence lines, right-of-ways, roadsides, canal banks, and levees. Seeds are spread primarily by birds and mammals through consumption and deposition of the fruit. Seeds are also spread by flowing water (Langeland and Burks 1998). Brazilian pepper seedlings will not tolerate inundation and are quickly killed; however large plants can withstand 6 months of flooding (Francis n.d.) with several feet of inundation. Brazilian pepper forms dense monocultures and completely shades out, crowds and displaces native vegetation. It also produces allelopathic agents that possibly suppress the growth of other plants. Brazilian pepper is a member of the family Anacardiaceae which includes plants such as poison ivy, poison oak, and poison sumac. The leaves, flowers, and fruits of Brazilian pepper produce a chemical that can irritate and form a rash on human skin and cause respiratory problems (Langeland and Burks 1998).

F.2.1.4 Cogongrass

Cogongrass (*Imperata cylindrica*) is a perennial grass that grows in compact bunches and produces extensive rhizomes. The leaf blades are erect and narrow with a whitish midvein off center and leaves can be one to four feet in length. The inflorescence is narrow, white, and plume-like. Cogongrass flowers in the spring, fall, and sometimes year-round. It produces seeds that are spread by wind, animals, and equipment. Cogongrass is native to Southeast Asia and was introduced into Florida in the 1930s and 1940s for forage and soil stabilization in Gainesville, Brooksville and Withlacoochee. More than 1,000 acres of cogongrass was established in central and northwest Florida by 1949. Cogongrass inhabits dry to moist sites and has been documented to occur in xeric hammocks, mesic flatwoods, herbaceous marshes, and floodplain forests (Langeland et al. 2008). It has extensively invaded disturbed areas such as fallow pastures (FDEP n.d.) and is commonly found along transportation and utility corridors (Langeland et al. 2008). Cogongrass forms dense stands which results in almost complete displacement of native plants. Dense stands of cogongrass also create a severe fire hazard, especially when mixed with other volatile fuels (FDEP n.d.).

F.2.1.5 Dwarf rotala

Dwarf rotala (*Rotala rotundifolia*) is a submersed aquatic plant and it is native to India and Southeast Asia. It was introduced into Florida's through the aquarium industry and was first found in a flood control canal in Broward County in 1996. Dwarf rotala is now inhabiting waters in Lee, Collier, Palm Beach, Broward, and Miami-Dade counties in Florida and Tuscaloosa County in Alabama (UF IFAS 2021/ EDDMapS). It is unique in its ability to grow fully submersed, emerged as well as terrestrially. Dwarf rotala roots at the nodes and grows year-round in Florida, which contributes to its invasive nature. This plant can reproduce from seeds and fragmentation. The growth habit of dwarf rotala leads to the creation of thick clumps that can block waterways and inhibit water movement and navigation. (SFER 2022)

F.2.1.6 Hydrilla

Hydrilla (*Hydrilla verticillata*) is a rooted submersed plant that is native to the Old World and Indo Pacific. It was likely introduced to Florida as an aquarium plant in the 1950's. Hydrilla can grow to the surface of waters as deep as 25ft and forms dense mats through the water column which displaces native plant communities. By the 1990s, hydrilla was widely distributed throughout the state, occupying more than 56,000 ha of lakes and rivers. Hydrilla supports the growth of a cyanobacterial epiphyte, *Aetokthonos hydrillicola*, which produces an avian toxin affecting herbivorous waterbirds and their avian predators (e.g., coots [*Fulica americana*] and bald eagles [*Haliaeetus leucocephalus*]; (Wilde 2005, 2014, Martin 2015).

F.2.1.7 Melaleuca

Melaleuca is an evergreen tree that can grow up to 100 feet tall. It has white flowers that form spikes often referred to as a "bottle brush." The fruit is a round woody capsule in clusters along the stem; each capsule can contain 200-330 tiny seeds. It is native to Australia and was introduced to Florida in 1906 as an ornamental plant and in the 1930s it was scattered over the Everglades in order to create forests (Langeland and Burks 1998). Melaleuca inhabits natural areas such as pine flatwoods, hardwood bottomlands, cypress forests, freshwater marshes, sawgrass prairies, and mangrove forests. It also infests disturbed sites such as improved pasture, natural rangeland, idle farmland, canal and levee banks and urban areas. It prefers sites that are seasonally wet. Melaleuca also flourishes in areas with standing water and persists in well-drained upland sites (Langeland and Burks 1998). Melaleuca displaces native plant species, reduces quality of wildlife habitat, alters fire regimes, and potentially alters wetland hydrology (Mazotti, Center, Dray, and Thayer 2008).

F.2.1.8 Old World Climbing Fern

Old World climbing fern (*Lygodium microphyllum*), is a plant that has long fronds that can grow up to 90 feet. The fronds grow along the ground, over shrubs or climb by twisting and winding around trees, vines and other structures. The rhizomes and rachis are wiry and they are brown to black in color. The leafy branches that form along the rachis are 2 to 5 inches in length and have many pairs of leaflets. It produces spores that are dispersed by the wind. In south Florida, the plant produces spores throughout the year. Each fertile leaflet of Old World climbing fern can produce up to 28,600 spores. Old World climbing fern is native to Africa, Asia, and Australia and the first record of it being found in Florida was in 1958. It was collected from a Delray Beach plant nursery where it was being cultivated (Langeland and Hutchinson 2005). Old World climbing fern has been documented to occur in hardwood hammocks, mesic flatwoods, forested swamps, wet flatwoods, hydric hammocks, floodplain forests, and strand swamps. It can completely overgrow the vegetation in these areas which allows the plant to compete with canopy trees and understory vegetation for light. The growth in the tree canopy provides an avenue for fire spread into the canopy which damages or even kills the trees. Over time, rhizomes accumulate in mats 3 feet or more thick on top of the soil (Langeland et al. 2008) which can prevent new growth of native plants. This plant is a threat to many areas within the project site (Ferriter et al. 2005) and disturbed sites.

F.2.1.9 Shoebuttan ardisia

Shoebuttan ardisia (*Ardisia elliptica*) is an evergreen, glabrous shrub or small tree approximately 17 feet tall. It was imported as an ornamental shrub as early as 1900 (Gordon and Thomas 1997). It invades understories of hammocks, tree islands, disturbed wetlands and cypress and mangrove areas.

This species often forms monocultures resulting in local displacement of native plant species. There is a tendency for reinvasion by shoebutton ardisia or other exotic plants following removal of dense thickets of this species. New infestations may go undetected due the physical similarity to the common native marlberry (*Ardisia escallonioides*).

F.2.1.10 Torpedograss

Torpedograss (*Panicum repens*) is a perennial grass that can grow up to 3 feet tall. It has extended rhizomes that can be rooted or floating. It has a panicle-type inflorescence that is 3-9 inches long. It flowers nearly year-round. Torpedo grass reproduces primarily through rhizome extension and fragmentation. It is native to Africa and Asia and was introduced into the Gulf Coast of the United States before 1876. Torpedo grass seed was introduced as a forage crop in the south and was planted in almost every southern Florida County by 1950. It is drought tolerant and grows in upland areas but thrives in areas with moist to wet sandy or organic soil. It inhabits scrub, coastal flatwoods, upper tidal marshes, mesic flatwoods, herbaceous wetlands, wet prairies, swales, lakeshores, canals, and other disturbed sites. Torpedo grass can quickly form a monoculture and displace native vegetation. In 1992, it was present in approximately 70% of the public waters in Florida. The largest population of torpedo grass was present in Lake Okeechobee. Approximately 14,000 acres of torpedo grass displaced native plants in Lake Okeechobee's marsh (Langeland et al. 2008). Torpedo grass is present in agricultural and water conveyance canals throughout the project area and has potential to spread into areas with the removal of levees and backfilling canals.

F.2.1.11 Water Hyacinth

Water hyacinth (*Eichhornia crassipes*) is a floating aquatic plant native to tropical South America that was introduced in Florida in 1884. The plant reproduces extremely quickly, it grows at explosive rates that exceed any other tested vascular plant (Wolverton and McDonald 1979). Vegetative reproduction occurs rapidly except in the coolest months. It forms large floating mats that block navigation, impact water control structures, degrade water quality, and dramatically alter native plant and animal communities (Gowanlock 1944, Penfound and Earle 1948). New plants are produced vegetatively and from seed, which germinate abundantly on exposed moist soils (Perez 2011). Water hyacinth has low nutrient needs and wide tolerance for water conditions that enables it to persistence and spread.

F.2.1.12 Water Lettuce

Water lettuce (*Pistia stratiotes*) is a floating aquatic plant native to South America. The plant reproduces extremely quickly except in the coolest months. It reproduces both vegetatively and from seed which are found to be up to 80% viable (Dray and Center 1989). Water lettuce was reported as early as 1765 by William Bartram as forming dense mats on the St. Johns River. It forms large floating mats that block navigation, impact water control structures, degrade water quality, and dramatically alter native plant and animal communities.

F.2.1.13 Water primrose

Many invasive aquatic *Ludwigia* species native to South and Central America have become widely established in Florida. These species include *L. grandiflora*, *L. hexapetala*, *L. peruviana*, and *L. peploides*. Water primrose initially spread horizontally across the water surface and crowd out native vegetation in the littoral zone of water bodies. When mature, emergent plants grow up to 4 to 5 feet

tall. In some waterbodies in Florida, creeping water primrose overwhelms populations of emergent native plants. Allelopathic effects further contribute to the plant's invasiveness (Dandelot et al. 2008).

F.2.1.14 Cattail

Cattails (*Typha spp.*) are native to Florida and occur in wetlands, lakes, rivers, canals, storm water treatment areas and other disturbed sites. Cattails grow up to 12 feet tall and have strap-like leaf blades. The inflorescence is spike-like with very tiny flowers. This plant is a primary native nuisance species within the project area. Many areas within the project area have been invaded by cattails. This is attributed to water with increased phosphorus being delivered to these areas which began in the late 1950s (Holmes et al. 2002). Areas where water control structures, conveyance features, and levees exist provide a suitable habitat for invasion and expansion of cattail.

F.2.1.15 Localized/Potential EDRR Species

Other non-native plant species of concern that are managed as an EDRR species for eradication include tropical American water grass (*Luziola subintegra*), mile-a-minute (*Mikania micrantha*) Tropical Nutrush (*Scleria macrocarpa*) and Wright's nutrush (*Scleria lacustris*).

F.2.1.15.1 Tropical American Watergrass

Tropical American watergrass (*Luziola subintegra*) is a perennial grass that is usually rooted but sometimes grows in floating mats. This plant can also grow in terrestrial sites. It produces a panicle type inflorescence. Tropical American watergrass spreads vegetatively and by seed. It is an aggressive grass that competes with both native and non-native invasive plants. It is native to Mexico, Central America, the Caribbean Basin with its range extending south through South America to Argentina (Krunzer and Bodle 2008). The first record of occurrence was in Lake Okeechobee in 2007 when two large mats of tropical American watergrass (approximately two and eighty hectares each) were found near Harney Pond Canal in Fisheating Bay at Lake Okeechobee. From the initial population identified, this plant quickly spread and by July 2009 more than 2,000 acres of the plant were treated in the lake. The plant was also found at the mouth of Fisheating Creek in both emergent and terrestrial forms. Since Fisheating Creek is the only unregulated flow into Lake Okeechobee, it is thought this area was the point of introduction. Since the initial sighting of tropical American watergrass in December 2007, other populations have been found in the Cody's Cove-Eagle Bay area, near Observation Shoal and inside Lake Okeechobee near the S-77 Structure and downstream in the Caloosahatchee River, C-43 canal. The majority of the populations of plants have occurred in areas that receive water flow from Fisheating Bay, however, one terrestrial population (in two small areas) was identified in the 8.5 Square Mile Area (SMA) adjacent to ENP. Through EDRR procedures, this plant was treated and appears to be eradicated in the 8.5 SMA. The source for the introduction of tropical American watergrass into 8.5 SMA is not known at this time, although contaminated equipment is highly suspected.

F.2.1.15.2 Mile-a-minute

Mile-a-minute (*Mikania micrantha*) is a federally listed noxious weed that recently appeared in south Florida. This vine has turned into a serious weed where it was introduced in Asia, Australia, and Africa (Holm et al. 1977, Zhang et al. 2004). Mile-a-minute was found near Homestead in 2008. An aggressive reconnaissance and eradication effort was launched immediately following the discovery of the plant. Controlling the plant is challenging due in part to plant populations on private lands (Dozier 2012),

although the threat of FDACS quarantine is an incentive for nursery owners to eliminate the weed. Eradication from Florida seems unlikely but containment and suppression remains a priority to prevent it from colonizing large natural areas like the South Dade Wetlands and ENP (SFER 2018).

F.2.1.15.3 Tropical Nutrush

Tropical nutrush (*Scleria macrocarpa*) is a perennial sedge and is native to the Tropical Americas, it occurs in several countries in Central and South America and the Caribbean. Although tropical nutrush has occurred in Florida since 2007, it was first identified in 2016 around Lake Hatchineha, in Polk and Osceola counties. It thrives along littoral areas and floodplains, forming dense, monospecific stands in the shaded understory of cypress and hardwood swamps. This species outcompetes and displaces native plant populations (Onisko 2020).

F.2.1.15.4 Wright's Nutrush

Wright's nutrush (*Scleria lacustris*) is a sedge native to seasonal wetlands of Africa, Central and South America, and the Caribbean. It has a robust, three-sided stem, with a side width up to 2.5 cm, and leaf length up to 30 cm. A red coloration can be seen at the base of the plant's stems. It was first recorded in Florida in 1988 in Jane Green Swamp, a marsh in the Upper St. Johns River Basin (UF/IFAS 2017). After range expansion, the current distribution of this species in Florida extends to more than twenty natural areas in seven counties (Brevard, Hendry, Indian River, Lee, Osceola, Okeechobee and Polk). In 2009, the Florida Exotic Pest Plant Council listed this plant as a Category I invasive species when recognition was made of its ability to alter the composition and structure of native wetland communities (UF/IFAS 2017).

F.2.2 Animals

Searches through existing data and resources indicate 65 animal species have been documented to occur within the project area. Other non-native animal species are probably present, however, documented citations could not be located. Information regarding species presence and distribution is largely incomplete for most taxonomic groups of animals. Not all of the 69 non-native animal species identified and documented to occur in the WERP area will have a significant impact on the ecosystem.

F.2.2.1 Widely Established Species

Species that are well established and are known or presumed to exert significant negative impacts on Florida ecosystems include the island applesnail (*Pomacea maculata*), purple swamphen (*Porphyrio porphyrio*), feral pig (*Sus scrofa*) Cuban tree frog, Asian swamp eel, and redbay ambrosia beetle (*Xyleborus glabratus*) and associated fungus (*Raffaelea lauricola*), veiled chameleon (*Chamaeleon calyptratus*), Mexican bromeliad weevil (*Metamasius callizona*), green iguana (*Iguana iguana*) and Burmese Python (*Python bivittatus*).

F.2.2.1.1 Redbay Ambrosia Beetle (laurel wilt)

Laurel wilt is a lethal disease of redbay (*Persea borbonia*) and other members of the Laurel family (*Lauraceae*). The disease is caused by a fungus (*Raffaelea lauricola*) that is introduced into trees by the wood-boring redbay ambrosia beetle (*Xyleborus glabratus*) (FDACS 2011). *Xyleborus glabratus* is the twelfth species of non-native ambrosia beetle known to have become established in the U. S. since 1990. All are suspected to have been introduced in solid wood packing materials, such as crates

and pallets (Haack 2003). Most native ambrosia beetles attack stressed, dead or dying woody plants, but *X. glabratus* attacks healthy Florida trees. Once infected, susceptible trees rapidly succumb to the pathogen and die. Besides redbay, it impacts other native and non-native members of the Lauraceae (Hanula et al. 2008) including swamp bay (*P. palustris*), an important species of many Everglades plant communities. Since its arrival in 2002, the red bay ambrosia beetle and laurel wilt have spread quickly throughout the southeastern U.S. In March 2010, the beetle was found in Miami-Dade County. Laurel wilt disease was subsequently confirmed on nearby swamp bay trees in February 2011. Aerial reconnaissance identified symptomatic swamp bay trees scattered throughout the Bird Drive Basin, northward into the Pennsuco Wetland area, and westward into ENP and WCA 3B. In February 2012, laurel wilt was also confirmed in the LNWR. There is currently no feasible method for controlling this pest or associated disease in natural areas. A systemic fungicide (propiconazole) can protect individual trees for up to one year, but widespread utilization in natural areas is impractical (Mayfield et al. 2009). State and federal agencies are monitoring the spread of laurel wilt disease and the red bay ambrosia beetle through the Cooperative Agricultural Pest Survey (CAPS) program. There is little to no research underway to assess the ecological impacts of laurel wilt disease. Interagency coordination is limited to the exchange of reporting information and some coordinated research. The red bay ambrosia beetle is considered a plant pest, so screening for additional introductions is carried out but is inadequate. Critical research areas include: (1) evaluating *Persea* resistance, (2) *Persea* seed/genetic conservation efforts, (3) potential chemical or biological control tools, (4) impacts on native plant communities, and (5) impacts on the Palamedes swallowtail butterfly (*Papilio palamedes*) and other host-specific herbivores.

F.2.2.1.2 Asian Swamp Eel

The Asian swamp eel (*Monopterus albus*) is a versatile animal, capable of living in extremely shallow water, traveling over land when necessary, and burrowing into mud to survive periods of drought (Shafland et al. 2010). This species is a generalist predator with a voracious appetite for invertebrates, frogs, and fishes (Hill and Watson 2007; Shafland et al. 2010). Wild populations in Florida originated as escapes or releases associated with aquaculture, the pet trade, or live food markets. Regional biologists are concerned that this species may become widely established, since the diverse wetland habitats of the Greater Everglades may be suitable for the species. Additionally, Asian swamp eels have a broad salinity tolerance giving concern that this species could also establish populations in estuaries (Schofield and Nico 2009). There are at least four reproducing populations of Asian swamp eels in Florida: North Miami canals, canal networks near Homestead adjacent to the ENP, eastern ENP, and in water bodies near Tampa (Collins et al. 2002; Nico et al 2011, USGS, personal communication, 2012; Jeff Kline, USNPS, personal communication, 2012). The impact of Asian swamp eels to Everglades fauna is undocumented and management options are currently limited to monitoring and electrofishing in canals. The species' generalist diet and adaptations to low water events suggests that native fishes, aquatic invertebrates, and frogs could be threatened. Nico et al. (2011) also report high parasitism rates in wild caught Asian swamp eels in Florida, raising concern that the species could be a vector for macroparasites to native fishes.

F.2.2.1.3 Cuban Treefrog

The Cuban treefrog is the largest species of treefrog in Florida and range from 1-4 inches in length. The Cuban treefrog has expanded pads on the ends of their toes which are exceptionally larger than toepads of Florida's native treefrogs. Cuban treefrogs have large eyes and usually have rough somewhat warty skin. Sometimes Cuban treefrogs have a pattern of large wavy marks or blotches on

their back and have stripes or bands on their legs. The color of the treefrogs varies from creamy white to light brown but Cuban treefrogs can be green, beige, yellow, dark brown or combination thereof. It is native to Cuba, the Cayman Islands, and the Bahamas. It was first reported in Florida in the 1920s in the Florida Keys, and was likely transported in cargo or ornamental plant shipments. Cuban treefrogs inhabit natural areas such as pine forests, hardwood hammocks and swamps. They also inhabit disturbed sites such as urban and suburban developments, agricultural areas such as orange groves and plant nurseries (Johnson 2007). Cuban treefrogs inhabit areas throughout most of the CERP area. These treefrogs are introduced to new areas as stowaways on cars, trucks, boat trailers and through shipment of ornamental plants and trees. Cuban treefrogs consume a variety of invertebrates and native treefrog species (Maskell et al. 2003). Native green and squirrel treefrogs (*Hyla cinerea* and *H. squirella*) are less likely to be found when Cuban treefrogs are present (Waddle et al. 2010), and when Cuban treefrogs are removed from an area, the abundance of native treefrogs increases (Rice et al. 2011). In addition, tadpoles of Cuban treefrogs are fierce competitors and can inhibit the growth and development of two species of native treefrogs (Johnson 2007). Effects of CERP projects on the distribution and abundance of Cuban treefrogs should be assessed given the Cuban treefrog's wide distribution and habitat tolerances, mounting evidence of direct impacts to native anuran species, and the lack of regional monitoring and control programs.

F.2.2.1.4 Feral Hog

Feral hogs (*Sus scrofa*), also known as wild pigs, have existed on the Florida landscape since their introduction four centuries ago. They are reported in all 67 Florida counties within a wide variety of habitats, but prefer oak-cabbage palm hammocks, freshwater marshes and sloughs and pine flatwoods. Although they do not favor marshes with deep water, during the dry season they make extensive use of partially dried out wetlands. Feral hog populations are particularly high in the counties immediately north and west of Lake Okeechobee, and in the Big Cypress and East Coast Regions. Hogs commonly grow 5-6 feet long with weights over 150 pounds. With a keen sense of smell and a powerful snout, they can detect and root up buried food. The diet of feral hogs includes vegetation, earthworms, insects, reptiles, frogs, bird eggs, rodents, small mammals, and carrion (Laycock 1966, Baber and Coblentz 1987). This invasive mammal is also known to prey on sea turtles, gopher tortoises, and other at-risk wildlife (Singer 2005). No animal native to North America creates the kind of disturbance when feeding that hogs do (Baber and Coblentz 1986). Rooting by feral hogs can convert native grassland and other low vegetation to what looks like plowed fields. Hog rooting may facilitate establishment of invasive plant species because invasive exotics typically favor disturbed areas and colonize more quickly than many native plants (Belden and Pelton 1975, Duever et al. 1986). Feral hogs are unusually prolific for large mammals. This is because they reach sexual maturity at an early age (6-10 months) (Barrett 1978), can farrow more than once a year (Springer 1977; Taylor et al. 1998), have large litters (4-8) (Sweeny et al. 2003), and often experience low natural mortality rates (Bieber and Ruf 2005). Recreational hunting is often a major source of mortality (Barrett and Pine 1980). In favorable habitat, however, hog populations are typically not greatly reduced by hunting (Bieber and Ruf 2005). There is no regional, coordinated monitoring program for the ubiquitous feral hog. Monitoring is limited to efforts associated with trapping programs and game management. Numerical monitoring of hogs present challenges because they are wary and adaptable animals that change their activity patterns and feeding areas in response to changing needs and threats from humans (Hughes 1985, Sweeny et al. 2003).

F.2.2.1.5 Green Iguana

The green iguana (*Iguana iguana*) is a large lizard native to Central and South America, extending to the eastern Caribbean (FWC 2018). Green iguanas can be found on the ground, in shrubs or in trees in a variety of habitats, from agricultural and natural areas to suburban developments. They are excellent swimmers, and are often found near canals and waterways. Male green iguanas can reach lengths of 1.5 meters, and can feed on a variety of vegetation, fruits, bird eggs, and dead animals. This species is characterized by its green coloration, a row of spikes down the center of the neck, back and upper section of the tail, which is banded with dark rings. Mature male iguanas display heavy jowls and a large throat fan, used both for sexual selection and self-defense. Green iguanas were first reported in Florida in the 1960s in Hialeah, Coral Gables and Key Biscayne along Miami-Dade's southeastern coast (FWC 2018). Breeding populations now extend along the Atlantic Coast in Collier and Lee Counties, and reports have been made as far north as Alachua, Highlands, Hillsborough, Indian River and St. Lucie Counties (FWC 2018). Those reported in more northern counties are likely individual pet releases, however, as green iguanas are not cold resistant, and will, therefore, be unlikely to establish breeding populations in these locations. In cleared habitats such as canal banks and vacant lots, green iguanas reside in burrows, culverts, drainage pipes and rock or debris piles. South Florida's extensive man-made canals serve as "ideal dispersal corridors to further allow iguanas to colonize new areas" (FWC 2018). Green iguanas cause damage to residential and commercial landscape foliage, and are often considered a nuisance by landowners. Some iguanas may even cause damage to infrastructure by digging burrows that erode and collapse sidewalks, foundations, seawalls, berms, levees and canal banks (FWC 2018). It is vital that this species be actively managed throughout South Florida to prevent further damage to infrastructure and native vegetation.

F.2.2.1.6 Purple Swamphen

The purple swamphen (*Porphyrio porphyrio*) is a member of the rail family native to Australia, Europe, Africa, and Asia. It is noticeably larger than its Florida native relatives, the American coot (*Fulica americana*), the common moorhen (*Gallinula chloropus*), and the purple gallinule (*Porphyrio martinica*). The swamphen and the gallinule both have purple plumage and red bills, but the face shield above the bill is red and the legs are pink in the swamphen while the face shield is pale blue, the legs are yellow and the bill has a yellow tip in the gallinule. Introduction of the swamphen was likely due to escapes from the Miami zoo and private aviculturists in Broward County. The purple swamphen feeds on shoots and reeds, invertebrates, small mollusks, fish, snakes, and the eggs and young of waterfowl (Pranty et al. 2000). Nests are typically large mounds of vegetation in wetlands. Known to be highly aggressive and territorial, the purple swamphen could negatively affect native water birds through competition for food and space and through direct predation. Rapid response efforts between 2006 and 2009 did not successfully reduce the abundance or distribution of this species. The management goal for the species has shifted from eradication to suppression (Jenny Ketterlin Eckles, FWC, personal communication, 2012). Efforts to remove birds by hunting did not significantly deplete the population. No other control tools are currently developed for this species. In recent years, purple swamphens have been sighted in the WCAs, ENP, Big Cypress National Preserve, Lake Okeechobee, and in all Everglades stormwater treatment areas. The FWC is currently conducting prey and habitat analyses to support a risk assessment, which will guide future management strategies (Jenny Ketterlin Eckles, FWC, personal communication, 2012). There are currently no coordinated monitoring efforts for purple swamphens.

F.2.2.1.7 Island Applesnail

The island applesnail (*Pomacea maculata*) is a large South American freshwater mollusk that is established throughout Florida. It was intentionally introduced through releases from aquaria and as a food crop. Potential impacts to the environment include destruction of native vegetation, competition with native fauna, and disease transmission. There is concern the island applesnail may out-compete the native applesnail, *P. paludosa* which is the primary food source of the endangered Everglade snail kite. In addition a newly described cyanobacterium (*Aetokthonos hydrillicola*) found in the Kissimmee Chain of Lakes is associated with a lethal neurologic disease, avian vacuolar myelinopathy (AVM), which affects avifauna in the southeastern United States (Wilde et al. 2005). Research has confirmed that bioaccumulation of a neurotoxin produced by *A. hydrillicola* in the island applesnail and birds fed with affected snail incur 100 percent development of AVM in laboratory birds (Dodd et al. 2016), suggesting a significant risk to the snail kite and other avifauna.

F.2.2.1.8 Burmese Python

Burmese pythons (*Python bivittatus*) are large (up to 5.5 meters) constrictors that are native to Southeast Asia (Dorcas et al. 2012) and are top predators (SFER 2013). For 20 years prior to being considered established, python sightings occurred intermittently in south Florida. In 2000, the Burmese python was considered established in south Florida and since that time, the population has increased significantly in abundance and geographic range (Dorcas et al. 2012). The Burmese python is found throughout the southern Everglades, particularly in ENP and adjacent lands including the East Coast Buffer lands and the northern ENP boundary along Tamiami Trail. Sightings have also been documented in the Key Largo region (SFER 2013). Pythons consume a wide variety of mammals and birds. More than 100 species have been identified as a food source and these include the endangered Key Largo woodrat (*Neotoma floridana smalli*) and the wood stork (*Mycteria americana*). In addition, American alligators (*Alligator mississippiensis*) are infrequently preyed upon by the python. Little is known about the impacts of predation by pythons on native species; however a recent study by Dorcas et al. indicates there has been a dramatic decline in mammal populations that coincides with the increase of pythons in ENP. The increase in the population size of pythons has been linked to a regional decline in small and medium mammals, but has not been distinguished from possible effects of changes in habitats and hydrology on mammal populations that also occurred during this time period (Dorcas et al. 2012).

F.2.2.1.9 Mexican Bromeliad Weevil

The Mexican bromeliad weevil (*Metamasius callizona*) was first detected in Florida in 1989. It was introduced via a shipment of bromeliads imported from Mexico. It is now found in many locations throughout south and central Florida (Frank and Cave 2005). The larvae of the weevil destroy bromeliads by mining into their stems. This insect is documented to attack 12 native bromeliad species. Ten of the species that are attacked by the weevil are state-listed as threatened or endangered and 1 of those species occurs naturally only in Florida. Two of these bromeliad species were listed due to the extensive damage incurred to their populations by the weevil.

F.2.2.1.10 Veiled Chameleon

The veiled chameleon (*Chamaeleo calyptrotus*) naturally occurs in mountain and coastal regions of the Arabian Peninsula, although it is also known to utilize a wide range of habitats. Florida populations of this species are suspected to have been established through intentional releases by reptile

enthusiasts. Breeding populations of the veiled chameleon are now documented in the Lee County (northwest estuaries), Miami-Dade County (one population near ENP a second adjacent to BCNP), Broward County, and Palm Beach County near the southern tip of LNWR (FWC 2013). In addition, reports of veiled chameleons are now common from Buckingham, Alva, Cape Coral, Marco Island, and Lutz, Florida. If chameleons continue to demonstrate the ability to spread from suburban and agricultural land and build populations in native Florida habitats, then the argument for an aggressive eradication program will be strong. (SFER 2018)

F.2.2.2 Localized/Early Detection Rapid Response Species

Of the species identified, there are four key carnivorous reptiles that are currently present within or in close proximity to the project area and have potential to cause significant ecological impacts. These include the Argentine black and white tegu, the Burmese python, northern African python, and the Nile monitor. At present time, these occurrences of the north African python, Argentine black and white tegu and the Nile monitor have been isolated but there is concern regarding further spread of these species from the southern portion of the project area. These reptiles are among south Florida's most threatening invasive animals. The species are considered top predators and increase additional pressures on native wildlife populations, particularly threatened and endangered species (SFER 2013). Other species considered EDRR include spectacled caiman (*Caiman crocodilus fuscus*), and the giant African land snail.

F.2.2.2.1 Northern African Python

Since 2001, over 40 northern African pythons (*Python sebae*) have been found in western Miami-Dade County (Jacob Kline, FWC, personal communication). This giant constrictor's natural history traits are similar to the Burmese python and is considered a high risk for establishment and expansion throughout South Florida (Reed and Rodda 2009). Rapid response efforts to eradicate this population are now of highest priority. The SFWMD, Miccosukee Tribe of Indians, and Miami-Dade County, the primary landowners within the Bird Drive Basin, are working closely with FWC and other agencies to address this threat.

F.2.2.2.2 Argentine Black and White Tegu

The Argentine black and white tegu is a large South American lizard that can reach 1.5 meters in length in the wild. Tegus seem to prefer savannas and other grassy open areas in its native range (SFER 2013). In Florida, tegus seem to prefer disturbed upland areas adjacent to wetlands or permanent bodies of water. These types of habitats are frequently found adjacent to canals and rock pits and occur throughout the South Florida landscape. Tegus are generalist predators with a diet that includes a variety of fruits, vertebrates, invertebrates and eggs. Because the tegu is a predator of eggs, it threatens native ground nesting birds and reptiles which includes threatened and endangered species such as the American crocodile (*Crocodylus acutus*) and Cape Sable seaside sparrow (*Ammodramus maritimus mirabilis*). Endangered snail species such as *Liguus fasciatus* are also potential prey. There are two known established populations in Florida, one in Hillsborough and Polk counties and one in southern Miami-Dade County. The population in Miami-Dade County seems to be increasing and expanding its range both to the west towards ENP and east toward Turkey Point. Both areas are home to endangered wildlife that may be threatened by tegus. Continued monitoring and removal efforts are needed to prevent the expansion into natural areas and control the population. Recently, there has been an increase in sightings near ENP which suggests the population is expanding. Systematic

surveys of the species are needed to validate the population is expanding near ENP (SFER 2011), and to provide early detection of possible range expansion to new areas.

F.2.2.2.3 Nile Monitor

The Nile monitor (*Varanus niloticus*) is a large, carnivorous lizard from sub-Saharan Africa that is capable of reaching 2.4 meters (FWC bioprofile). It is a generalist feeder and an egg specialist in its native range (SFER 2013) that will feed on a wide variety of invertebrates and vertebrates it acquires by either predation or scavenging (FWC bioprofile). As such, the Nile monitor could impact a variety of native and threatened species in Florida through both competition and predation. The Nile monitor may pose a serious threat to a number of wading birds, marsh birds, gopher tortoises (*Gopherus polyphemus*), burrowing owls (*Athene spp.*), Florida gopher frogs (*Lithobates capito*), sea turtles and other ground nesting species. They may negatively impact populations of American alligators and American crocodiles via egg predation and competition (FWC bioprofile). The Nile monitor has been well established in the Cape Coral area since the 1990s. There is also a small breeding population near Homestead Air Force base in Miami Dade County (SFER 2011). More recently, a breeding population of Nile monitors has been discovered in Palm Beach County and numerous reports of the species throughout Broward County also suggest a breeding population. Because of their threat to our native wildlife, this species has potential to impact restoration efforts.

F.2.2.2.4 Spectacled Caiman

The spectacled caiman (*Caiman crocodilus fuscus*) naturally occurs throughout Central and South America, and can reach sizes of about 2.4 meters. They are easily distinguished from native crocodilians not only by their smaller adult size, but by the characteristic vertical dark bands that can be found on their tails. In Florida, spectacled caiman are commonly encountered in ditches, canals, and disturbed wetlands but are occasionally found in relatively undisturbed marshes. This species was first reported within canals at the Homestead Air Force Base as early as 1960 (Ellis 1980). It feeds primarily on fish, mammals, waterbirds, and snails in its native range (Thorbjarnarson 1993). Breeding populations are documented in localized areas of Miami-Dade and Broward counties. Given its intolerance of cold temperatures, breeding populations will remain limited to southern Florida. (SFER 2018)

F.2.2.2.5 Giant African Land Snail

The Giant African land snail (*Lissachatina fulica*) is a large snail native to Africa, but was discovered in Miami in 2011 (USDA 2013). It is known to eat a variety of vegetation, namely crop plants, horticultural plants and environmentally valuable plants. This species of snail is an intermediate host of the rat lungworm (*Angiostrongylus cantonensis*), which can spread meningitis to humans (Cowie 2013). This lungworm was undetected in Florida prior to the Giant African land snail's introduction. A previous infestation of this snail occurred in Miami in 1966, and the State of Florida spent \$1 million and 10 years of effort on eradication (USDA 2013). The Giant African land snail is known to occur in developed areas of Broward and Miami-Dade counties, from Davie south to Homestead. As of July 2017, researchers have identified 31 population cores in Miami-Dade County and a single core in southern Broward County. There are indications that control efforts are having an effect, as fewer large snails are being reported, and local eradications of the snail are being observed in some of the population cores (Roda et al. 2016).

F.3 Introduction to Invasive Species Management

Invasive species management includes prevention, monitoring, education and public awareness, EDRR, control and management as well as adaptive management. In addition to these components, it is important to understand the risks and uncertainties associated with invasive species in order to effectively implement control/management measures and to adaptively manage.

F.3.1 Prevention

Prevention is the first line of defense and the most efficient and cost-effective approach to reduce the threat of invasive non-native species. Successful prevention will reduce the rate of introduction and establishment and thereby reduce the impacts of invasive species. One essential element to prevention is identifying the high-risk pathways that facilitate introductions and implementing actions to impede those introductions. Other critical elements include using effective management tools to reduce unintentional introductions and using risk assessment for both intentional and accidental introductions of non-native species. Baseline data and monitoring systems are required in order to evaluate the success of preventative measures.

F.3.2 Education and Public Awareness

A key to addressing problems caused by invasive species is to increase public awareness of their impacts and providing information about how individuals can help prevent the introduction and spread. However, reaching each person whose activities may affect our natural environment is a daunting task. Collaboration, cooperation and coordination across federal and state agencies, local governments, tribal entities, and the public and private sectors is required to facilitate this effort.

F.3.3 Monitoring

Natural resource managers need spatial data on invasive species populations to develop management strategies for established populations, direct rapid response efforts for new introductions, and evaluate the success of control efforts (Myers et al. 2000; Dewey and Andersen 2004; Barnett et al. 2007). Several approaches may be taken to document the spatial distribution and population trends of invasive species. Each method has strengths and weaknesses and should be utilized according to specific management objectives. Monitoring is the collection and analysis of population measurements in order to determine changes in population status and progress towards meeting a management objective (Elzinga et al. 1998). This type of monitoring is usually intended to detect relatively small changes in populations over time and often utilize small scale plots and/or transects. Invasive species surveys and inventories may be preferred when the objective is to detect populations and describe their spatial distributions over large landscapes, especially when early detection of new populations is desired (see EDRR discussion below).

Optimally invasive plant mapping methods have high positional accuracy, high species detection accuracy (particularly for low-density infestations), rapid turnaround time, relatively low cost, and the ability to quantify the degree of infestation (USDA 2012). Ground-based surveys can provide high positional accuracy and species detection, but can be time consuming and logistically unrealistic for large landscapes (Rew et al. 2005). Stratified subsampling approaches to ground surveys can mitigate some of these limitations but probabilistic mapping may be ineffective for early detection needs of land managers (Barnett et al. 2007) and may not provide sufficient fine scale information over large areas.

Developments in remote sensing technology have greatly improved opportunities for rapidly obtaining spatially precise data on invasive plant populations, particularly for large areas (Lass et al. 2005). However, the ability to detect target species using remote sensing is still limited to conditions where the species has a unique spectral signature or is a dominant canopy species and is often ineffective at detecting target species at low densities (Shafii et al. 2003). This inability to detect target species at low densities is a significant limitation for land managers focused on containment of expanding populations and detection of new invasions. Visual surveys from aircraft have been effectively used to map invasive plant distributions in the Everglades since 2008 (Rodgers et al. 2014). While visual aerial surveys may provide cost-effective information on landscape distributions of targeted plants, it has limited value for long-term change detection or fine scale assessments of abundance. This method may also lack sufficient detection precision for small plant species or species that occupy understories. Use of UAVs may also provide relatively inexpensive invasive plant monitoring data and video documentation provides a permanent record of conditions. However, detection accuracy may be less than that of visual surveys, especially at low densities or new species introductions.

F.3.4 Early Detection and Rapid Response

Once a species becomes widespread, the cost to control it will more than likely require significant and sustained funding. EDRR may be a cost-effective strategy to locate, contain, and eradicate invasive species early in the invasion process in order to minimize ecological and economic impacts of non-indigenous species (Rejmanek and Pitcairn 2002).

The three components of EDRR are *Early Detection*, *Rapid Assessment*, and *Rapid Response*. Early detection is defined as a comprehensive and integrated system of active or passive surveys to locate, identify, and report new invasive species as quickly as possible in order to implement procedures when it is feasible and less costly. Rapid Assessment includes the actions necessary to determine the appropriate response. This assessment identifies the current and potential range of the infestation, an analysis of the risks associated with the invasion, and timing and overall strategy for the appropriate actions. Rapid response is defined as a systematic approach to control, contain, or eradicate these species while the infestation is still contained in a particular area. Based on the results of the rapid assessment, a rapid response may be implemented to address new introductions or isolated infestations of a previously established species invading a new site (i.e., containment strategy).

Another critical element to rapid response is having the infrastructure in place to quickly implement management actions while new invasions can still be eradicated or contained. Effectively implementing EDRR will require coordination and collaboration among federal, tribal, state, local governments, nongovernment organizations (NGOs) and the private sector (National Invasive Species Council 2008).

F.3.5 Control and Management

Integrated Pest Management (IPM) is an effective approach to manage invasive species. IPM is the coordinated use of the most appropriate strategy to prevent or reduce unacceptable levels of invasive species and their damage by utilizing the most economical means, and with the least possible hazard to people, property and the environment. Physical, mechanical, chemical and biological control methods are utilized in IPM.

Physical control, sometimes referred to as cultural control, is the physical manipulation of an invasive species or their habitat. A number of techniques are used for physical control. These include manual removal, installing barriers and environmental alterations such as water level manipulation, prescribed fire, and light attenuation.

Mechanical control refers to the use of machinery designed to cut, shear, shred, uproot, grind, transport and remove invasive species. Equipment used to complete mechanical control may include but is not limited to heavy equipment such as an excavator or front-end loader (with a root rake, grinding heads or other attachments), cutter boats, dredges and mechanical harvesters (Haller 2009).

Chemical control is the use of a specially formulated pesticide to control an invasive species. The United States Environmental Protection Agency defines a pesticide as “a substance or mixture of substances intended for the prevention, destruction, repulsion, or mitigation of any pest”. The term pesticide encompasses a broad range of substances including herbicides, insecticides, fungicides etc. Pesticides are applied through ground and aerial applications.

Biological control, also known as bio-control, is the planned use of one organism to suppress the growth of another. Biological control is primarily the search for and purposeful introduction of species-specific organisms that selectively attack a single target species. Organisms such as insects, animals, or pathogens that cause plant diseases are used as biological controls (Cuda 2009).

Objectives of management can include complete eradication within a given area, population suppression, limiting spread and reducing effects of invasive species. Once an invasive species becomes widely established complete eradication is usually not feasible. The most effective action for managing widely spread invasive species is often preventing the spread and reducing the impacts by implementing control measures. This concept is known as maintenance control. Maintenance control is defined as controlling an invasive species in order to maintain the population at the lowest feasible level.

F.3.6 Risk and Uncertainties Related to Invasive Species

As with most land management activities, there are a number of risks and uncertainties associated with invasive species management. The use of an adaptive management approach will help develop and prioritize invasive species control strategies. As restoration proceeds, invasive species may establish and/or spread as a direct result or independently of restoration activities. In the context of LRWRP and the long-term management of the natural resources within the study area, risks include but are not limited to:

- Introduction of new invasive species which are difficult to control and/or new species for which techniques are unknown or haven't been developed. Restoration activities which unintentionally facilitate the spread of invasive species via contaminated earth moving equipment.
- Undetected spread of invasive species into new areas, making containment of populations more costly and less likely to succeed.
- Uncontrolled invasive species which create disturbances or alter ecosystems such that desired restoration outcomes are not achieved.
- Failure to secure necessary funding to control invasive species.

- Undesirable impacts on non-target species and ecosystem functions resulting from invasive species control efforts.
- Not taking action to manage a species due to inaccurate assessments of the species impact on restoration activities.

The major uncertainty is that in most cases we do not have necessary information for detailed, specific pre-project evaluations of the need for management activities to control invasive species. With the exception of a few well-established and well-studied species (e.g., melaleuca), there is an information deficit on the status, potential impact, and effective control techniques for priority species. This is particularly true for non-indigenous animals. Current knowledge on invasion mechanisms suggests that some restoration activities may facilitate the spread of certain priority species. For example, partial removal of canals and levees could encourage spread of or provide sites for colonization by numerous invasive species, including Brazilian pepper, Old World climbing fern, Nile monitors, pythons, and Cuban treefrogs. However, there remains considerable uncertainty regarding the degree to which different species will respond, if at all, to restoration activities and how these responses will impact achievement of restoration goals.

Given the high degree of uncertainty, the most effective and lowest cost management option is early detection and rapid removal of invasive species during and post project. Central to this strategy is the implementation of a rigorous monitoring program (discussed below).

One specific uncertainty has been identified in the initial analysis of the selected plan. It is listed here to provide a starting point for developing monitoring, control and BMP strategies for the construction and operations phases of the restoration. This uncertainty is addressed by the WERP Adaptive Management Plan as well.

- WERP AM Uncertainty ID#5: What is the appropriate methodology to use in restoring vegetation (i.e. mechanical means, herbicide, fire) in areas identified as “wetland restoration”? (Objective 1) (The areas identified as wetland restoration are infested with invasive non-native species.

F.4 Existing Management Programs

Management of invasive species within the project area is conducted by several agencies. The magnitude of the control programs within the project area is dependent upon the level of funding available. Portions of allocated funding for these programs have been and potentially will be redirected to other programs in the future. Management activities vary in effectiveness which also influences species control and spread within the project area.

F.4.1 South Florida Water Management District

The SFWMD manages invasive exotic aquatic and terrestrial plants in canals and on levees within the project area, interim project lands, and on public conservation lands. Most of the vegetation management is outsourced through the Vegetation Management Division and includes herbicide application contractors, mechanical removal contractors, and use of biological controls such as plant specific insects and herbivorous fish. The Melaleuca Control Program is a major focus for the SFWMD, but other priority plant species are controlled within the WERP study area as funding resources allow.

F.4.2 US Army Corps of Engineers

The USACE also conducts treatments of priority species on the Herbert Hoover Dike. In addition to the operations and maintenance program on Lake Okeechobee, the USACE conducts treatments of vegetation during the construction & OMRRR phase for CERP projects. Vegetation treated includes FLEPPC Category I and II species, as well as native nuisance species.

F.4.3 U.S. Department of Agriculture / University of Florida

The SFWMD, USACE, NPS, USFWS, FWC, and other agencies provide financial support to the U.S. Department of Agriculture – Agricultural Research Service (USDA-ARS) and the University of Florida (UF) for the development of invasive plant biological controls. Efforts to identify safe and effective biological controls have led to important advancements in the integrative management of several invaders, including melaleuca, Old World climbing fern, water hyacinth, and alligator weed. The *CERP Melaleuca Eradication and Other Exotic Plants – Implement Biological Controls Project* is dedicated to the implementation of biological control agents once overseas surveys and quarantine testing has developed agents deemed safe for release in Florida. The project includes a mass rearing annex to the existing USDA-ARS biological control facility in Davie, Florida, in support of implementing the mass rearing, field release, establishment, and field monitoring of approved biological control agents for melaleuca and other invasive nonindigenous species.

F.4.4 Florida Fish and Wildlife Conservation Commission

The FWC's Invasive Plant Management Section is the designated lead entity in Florida responsible for coordinating and funding the statewide control of invasive aquatic and upland plants in public waterways and on public conservation land. In addition to funding the SFWMD melaleuca control program, FWC annually awards funding for individual invasive plant management projects in the Everglades region. Allocation of control funding is determined by an interagency regional working group.

F.4.5 Invasive Animals

Efforts to develop control tools and management strategies for several priority species are underway for a few priority animal species. These include the Burmese python and other giant constrictors, the Nile monitor, and the Argentine black and white tegu. Control tools are very limited for free-ranging reptiles, and the application of developed methods is often impracticable in sensitive environments where impacts to non-target species are unacceptable. Available tools for removing large constrictor snakes and lizards currently include trapping, detection dogs, and visual searching. Potential tools include the use of toxicants, introduced predators, and pheromone attractants, but these have not been fully explored to date.

Regional biologists have developed a conceptual response framework for established priority invasive animals in south Florida. Objectives within this framework are classified into three main categories—containment (slow the spread), eradicating incipient populations (remove outliers), and suppression (reduce impact in established areas). The resources to implement this strategic framework remain insufficient, but close collaboration between agencies has allowed for some coordinated efforts. Currently, FWC, NPS, UF, and SFWMD are conducting trapping and visual searching for Burmese pythons, northern African pythons, Argentine black and white tegus, spectacled caimans, and Nile monitors.

F.5 Existing Monitoring Programs

Since 2008, the SFWMD and USNPS, along with other partner agencies, have utilized digital aerial sketch mapping (DASM) for a region-wide mapping program over 728,000 ha in the Everglades. DASM is a method for mapping plant infestations “on-the-fly” using GPS-linked computers and trained biologists. Visual surveys allow an observer to learn to recognize targeted species, sometimes at low densities, under a range of environmental and phenological conditions. Visual aerial surveys also may provide data more rapidly than other methods, which is important when rapid responses to newly established threats are expected. The primary objective of the DASM inventory program is to determine the distributions of four priority invasive plant species on managed conservation lands in the region. These are Australian pine, Brazilian pepper, melaleuca, and Old World climbing fern. A secondary objective of the program is to detect new plant species invasions in remote areas to facilitate rapid response efforts. This data is currently collected on a two-year cycle.

Since 2010, the SFWMD has been collaborating with UF, FWC, USGS, NPS and FWS on the Everglades Invasive Reptile, Amphibian, and Mammal Monitoring Program (EIRAMMP). The purpose of the project is to develop an early detection, rapid response, removal and monitoring program for invasive reptiles and amphibians within Greater Everglades ecosystems. Specifically, the program seeks to (1) determine the status and spread of existing populations and the occurrence of new populations of invasive reptiles and amphibians, (2) provide additional EDRR capability for removal of invasive reptiles and amphibians, and (3) evaluate the status and trends of populations in native reptiles, amphibians, and mammals. The monitoring program involves visual searches for targeted invasive species on fixed routes along levees and roads within LNWR, WCA-2, WCA-3, Big Cypress National Preserve (BCNP), Southern Glades Wildlife Management Area, ENP, Corkscrew Swamp Sanctuary, and other areas such as the C-51 canal, US Highway 1, and Card Sound Road. Visual searches and call surveys are conducted to monitor invasive species and their potential prey species. Twenty-one routes have been established, and seven are active. The encounter rates for Burmese pythons ranged from 0.0014 to 0.0035 observations per kilometer. To date, a total of 105 Burmese pythons have been detected during these visual surveys. In 2018, EIRAMMP will increase focus on removal of this priority species.

F.6 Management Strategy and Plan

WERP includes the construction of two STA’s with associated discharge and flow way infrastructure, the plugging of existing canals, the installation of levee tie-ins, the degradation of existing levees, the backfilling of existing canals, the restoration of areas impacted by nuisance and non-native invasive vegetation, the restoration of historic tree islands bisected by the C&SF Project infrastructure, and the installation of water management structures (gated culverts, ungated culverts, weirs, etc.).

Many of the new infrastructure features and created flow ways, as well as construction and operations and maintenance activities, have the potential to spread and promote establishment of non-native invasive and native nuisance species. Proposed restoration activities may affect ecosystem drivers that directly or indirectly influence the invasiveness of non-native species. These factors may affect invasive species positively or negatively, depending on the unique characteristics of individual species and the environmental conditions for a given biological invasion (Doren et al. 2009). Many of the areas where features are proposed are currently inhabited by non-native invasive and native nuisance species. Construction of the proposed features has the potential to spread the existing non-native invasive and native nuisance species on site as well as introduce new invasive species via

contaminated equipment. Disturbed areas resulting from construction are likely to become established with non-native invasive and native nuisance species. New flows created by operations of the proposed features may serve as vectors to spread invasive and native nuisance species into new areas. Monitoring is a critical component of the management strategy. Information on distribution and restoration responses of invasive species should be used to inform decisions on control strategies. Invasive species surveillance, monitoring, and control should be carried out within the construction footprints, as well as impacted areas. Species of non-native vegetation to be treated include, but are not limited to, species listed in the current version of the FLEPPC invasive plant lists and the Florida Department of Environmental Protection prohibited plant list. The priorities for managing vegetation include FLEPPC category I and II species, new invasive plant introductions, native nuisance species and plants that impact project operations. Management of animal species will include surveillance, control, and monitoring.

The strategy for managing invasive species will be to utilize an IPM approach. Objectives of management will include complete eradication, population suppression, limiting spread and reducing effects of invasive species. Eradication will be the objective for new established species that are localized. The objective for wide-spread invasive species will be to implement control measures to suppress and prevent the spread of identified priority invasive species.

F.6.1 Surveillance – Early Detection and Rapid Response

EDRR should be implemented during every phase, for the life of a project. EDRR is an effective management measure to controlling and containing invasive species that were not previously within the project area. EDRR minimizes the negative impacts the invasive species has on the ecosystem and economy and reduces future treatment and management costs. It is very difficult to predict when and where an invasive species may appear. As such, estimating a needed budget is near impossible. However, to assist managers, a priority list of species to immediately respond to under EDRR management strategy has been developed.

A framework for establishing an EDRR program in the Everglades was recently drafted by an interagency team of invasive species experts and land managers (see ECISMA EDRR Plan at http://www.evergladescisma.org/ECISMA_EDRRPlan_2009-2011.pdf). As discussed above (Section F.3.4 Early Detection and Rapid Response), EDRR includes three strategy elements: 1) early detection, 2) rapid assessment, and 3) rapid response.

1.) Early Detection: This plan proposes implementation of routine surveillance in the project area in order to minimize the time between initial introduction and detection of a new species. Strategic surveillance by trained biologists in proximity to the CEPP project elements should greatly increase the probability of detection of new species. In many cases, existing programs could be expanded to include focused monitoring in the CEPP footprint. For example, the EIRAMMP is well suited for enhanced surveillance for numerous invasive animal species (see Section F.4 Existing Management Programs).

2.) Rapid Assessment: Following the detection of new invasions (or expansion of formerly contained invasions), it is important to gather and process available information to determine the potential risk and control options in the face of high uncertainty. Critical questions must be answered in a relatively short period of time. Example questions include:

- What is the spatial extent and abundance of the invasive non-native species?

- What is the likelihood that the species will impact native species, ecosystem function, operations infrastructure, or human health?
- What are the management options for containment or eradication?

Numerous tools are available to assist natural resource managers with the assessment phase of EDRR, though none of them is likely to be 100% accurate in assessing the risk of a species. This plan proposes utilization of the IFAS Assessment of Non-native plants in Florida's Natural Areas, the Florida Exotic Pest Plant Council's Invasive Species List, the FWC Non-native Animal Bioprofile protocol, and the ECISMA Rapid Response Plan for assessing the risks of non-indigenous species in the CEPP footprint. These assessments should be conducted with CEPP biologists, subject matter experts, and stakeholders.

3.) Rapid Response: This is the "risk management" component of EDRR. Once a species is determined to have a high probability of ecological impact and control options are available, rapid response strategies aimed at containment, and ultimately eradication, can be formulated and implemented. To be effective, rapid response programs must have built in procedural, financial and logistical capacity to respond quickly to newly established threats. Since it is not possible to accurately predict the number and severity of new invasions during the project, this plan proposes contingency funding for rapid response activities in the event new, high-priority species establish in the project area. During the pre-construction phase, protocols for implementing rapid response should be developed.

F.6.2 Control

A combination of biological, physical, mechanical, and chemical control methods will be utilized to manage invasive species.

Biological control agents will be used to decrease the targeted invasive species competitive advantages over native species and to weaken the invading population by increasing leaf mortality, decreasing plant size, reducing flower and seed production, and/or limiting population expansion. Biological control agents will be acquired through the "Melaleuca Eradication and Other Exotic Plants – Implement Biological Controls" project, which is a component of CERP. One element of this CERP component includes the implementation of biological control agents which involves mass rearing, field release, establishment and monitoring of approved biological controls in south Florida and the Everglades. The four main invasive plant species targeted for control through this component include melaleuca, Australian pine, Brazilian pepper and Old-World climbing fern.

It is anticipated that physical control methods will be limited. Prescribed burns will be conducted in order to promote native plant growth and should be planned, if possible, to target invasive species when they are most susceptible to fire. Hand pulling of melaleuca and other non-native plant species will occur when it is feasible. Weed/debris barriers will be placed at water control structures when it is required to minimize dispersal of floating vegetation. Physical control measures will be utilized for invasive animal control. Examples of these measures include trapping of feral hogs, controlled harvest/overfishing (nets, fishing tournaments specific to invasive fish species) and compliance with FWC Fishing Regulation release/movement of fish (no return to water/used as bait).

Mechanical control will be implemented to remove non-native plant species when the construction of project features requires such removal. Heavy equipment such as bulldozers, front-end loaders and excavators (with or without grinding heads) will be utilized to uproot, grind and/or clear and grub. It

is expected this type of control method will be utilized during levee degradés, canal backfilling and during construction of new project features such as water control structures.

Chemical control will be utilized to treat aquatic and terrestrial invasive plants. Methods for treatment will include hack-n-squirt, basal bark, cut-stump, foliar and aerial application. EPA approved herbicides will be utilized to control invasive plants. Chemical control will be utilized to treat invasive plants in canals, along levees and in wetland/natural areas etc.

F.6.3 Monitoring

Monitoring of invasive species populations will be conducted through DASM, Unmanned Aircraft System (UAS) surveys, electrofishing and EIRAMMP. Invasive species will also be identified through monitoring for the Adaptive Management Plan. This information will be provided to invasive species managers to ensure appropriate management measures are implemented.

F.6.4 Pre-construction Phase

Baseline conditions need to be established prior to the construction phase. Existing monitoring programs should be used as much as possible to establish baseline conditions prior to construction activities beginning. Although there are no system-wide monitoring programs for invasive species in the Everglades region, several individual agencies collect data. Data mining will be the primary resource to obtain baseline data via collaboration with the individual agencies. In areas with data gaps, surveys will need to be accomplished by the most cost-effective method (e.g. ground survey, Unmanned Aircraft Systems survey, DASM).

Existing monitoring and management programs should continue to be implemented. The existing programs help maintain invasive and nuisance species at a controlled level.

A significant length of time lapses from the time a project is planned to when it receives congressional authorization and appropriations, and ultimately goes to construction. As property (lands and structures) sit with no activity, vegetation, and wildlife changes can occur. Unmanaged areas become inhabited by many species of flora and fauna, native and non-native. Older growth vegetation is more difficult and more costly to treat / remove versus lands that are managed along the way. As these lands become established with invasive species, there is an increased risk of spreading the invasive species to neighboring lands. Therefore, it is beneficial, ecologically and economically, to manage the lands early on. Managing invasive vegetation throughout the interim phase reduces construction costs since mowing is much less costly than clearing/grubbing and treating, and rapid response of new infestations helps reduce spread into environmentally sensitive areas. Site 1 Impoundment is an excellent example. \$2.9M is estimated to manage invasive species during construction and until turnover to the local sponsor. The property's prior use included plant nurseries and pasture. Once project lands were acquired by the sponsor, the land sat unused until the Site 1 project was ready to begin construction. By this time, the project lands became highly vegetated, primarily by invasive species. It would have been significantly less expensive to have maintained the lands until the time of construction versus waiting until construction started.

F.6.5 Design and Construction Phases

The best method of controlling invasive and nuisance species is to prevent non-native species from being introduced and established to begin with. Incorporation of invasive species prevention and

control into project designs, alternatives analysis, and operational plans has the potential to save significant resources during the long-term. The plans and specifications phase should simply design “with the end in mind.” When the end goal is ecosystem restoration, the designers should periodically obtain input from invasive species experts to identify design features and operation strategies that could potentially favor the establishment and spread of invasive species. An example of design influences on invasive species is levee removal without backfill of canals. Without canal backfilling, deep water refuges for non-native fishes and invertebrates (from both seasonal cold temperatures and seasonal drying) are maintained, and barriers to dispersal from canal waters to marsh habitats are removed. Design alternatives should be explored that would allow seasonal cooling of water in the canals. Cooler water temperatures will reduce the refuge capacity for cold temperature sensitive non-native fishes. In some cases, such as the coastal canals, aquatic barrier technologies could be used to mitigate the spread of non-native aquatic species.

Below are examples of cost-saving measures to consider during design and construction.

- Include invasive species management staff from the Corps, SFWMD, and other partner agencies throughout the design and construction phases.
- Work with subject matter experts to identify design features that may create habitat or entry points for invaders. Evaluate design alternatives to mitigate potential design vulnerabilities.
- Design to promote the establishment of native species.
- Use construction methods that minimize ground disturbance whenever possible.
- Contain mobilized nutrients resulting from soil disturbances.
- Require all construction contractors to follow vehicle and equipment decontamination protocols prior to deployment. Coordinate with invasive species specialists for decontamination protocol specifications.
- Evaluate cost/benefit ratios for treating invasive/nuisance species prior to construction activities. In some cases, pre-construction removal of a species may significantly reduce its spread.
- Implement a monitoring and rapid response protocol aimed at detecting and controlling new invasions early.
- Manage and control invasive/nuisance species during the entire construction phase.
- When native planting is specified in the plans, use plant material from regional sources that are weed and pathogen free.

Construction will be the responsibility of either the Corps or the SFWMD. This will be determined at a future time. Regardless of which agency will be responsible, both agencies commit to requiring the construction contractor to implement preventive measures and best management practices that will minimize the potential introduction and spread of invasive and nuisance species due to construction equipment (including personal protective equipment) and activities. This commitment is also included in the Project Implementation Report/Environmental Impact Assessment (Section 5.2.5 Environmental Commitments).

The Corps currently includes the following language in all of their specifications (Specification # 01 57 20 Environmental Protection, “Prevention of Invasive and Nuisance Species Transfer”):

The Contractor shall thoroughly clean equipment prior to and following work on the project site to ensure that items/materials including, but not limited to, soil, vegetative debris, eggs, mollusk larvae, seeds, and vegetative propagules are not transported from a previous work location to this project site, nor transported from this project site to another location. Prevention protocols require cleaning all equipment surfaces, including but not limited to, undercarriages, tires, and sheet metal. All equipment, including but not limited to, heavy equipment, vehicles, trailers, ATV's, and chippers must be cleaned. Smaller equipment, including, but not limited to, chainsaws, loppers, shovels, and backpack sprayers, must be cleaned and inspected to ensure they are free of eggs, vegetative debris, vegetative propagules, etc. The Contractor may utilize any method accepted by the Government; common accepted methods include pressure washing and steam cleaning/washing equipment. Prevention protocols should also address clothing and personal protective equipment.

Prior to the commencement of work, the Contractor shall complete and provide an invasive and nuisance species transfer prevention plan to the Corps for approval. This plan shall be part of the Environmental Protection Plan as defined in subparagraph "Environmental Protection Plan" of paragraph SUBMITTALS (Part 1.5) above. The invasive and nuisance species transfer prevention plan shall identify specific transfer prevention procedures and designated cleaning sites/locations. Prevention protocols may vary depending upon the nature of the project site. It will be the responsibility of the Contractor to ensure all equipment coming onto and leaving the project site is inspected and not harboring materials that would spread, or potentially spread, invasive and nuisance species onto or off the project site. The Contractor shall provide a report verifying equipment brought on site was cleaned and shall provide a report verifying equipment was cleaned prior to removal from the project site.

F.6.6 Operational Testing and Monitoring Period

The operational testing and monitoring period is the timeframe from the end of construction until the project is transferred and accepted by the local sponsor. EDRR is very critical and the most cost-effective management measure during this period. Disturbed areas, such as areas impacted from construction activities, are prone to the establishment of invasive and nuisance species. Early detection of invasive and nuisance species and immediate treatment/control measures prevent these species from establishing and becoming long-term problems, ecologically and economically.

F.6.7 OMRR&R Phase

"Prevention of Invasive and Nuisance Species Transfer" language applies not only to the construction phase, but also to the OMRR&R phase. The preventive measure applies to contractors and government employees. Maintenance equipment and rental equipment are often used at multiple locations. As equipment is moved from one location to another, this potential spread vector can easily be reduced / prevented simply by ensuring the equipment is clean prior to arrival on site and prior to leaving the site.

In addition, numerous operational aspects of the restoration can influence mechanisms of invasion. For example, many non-indigenous species become more invasive in environments with elevated nutrient availability. With large pulses of only slightly elevated phosphorus levels, some invasive plant species could establish and spread.

F.6.8 Specific Control by Project Feature – Construction Phase

Surveillance and management of invasive species may begin as soon as Construction Phase is initiated. This will be in effort to minimize spread of priority species during the construction phase. Various management measures will be implemented in order to reduce colonization and spread of invasive plant and animal species. Once management has been completed, further measures shall be implemented during the construction phase in order to prevent re-establishment of populations of invasive species.

F.6.8.1 Region 1

North Feeder STA

The NFSTA includes 3,700 acres of effective treatment area. Features of the STA will include the north feeder intake Canal, distribution canal, outflow canal, two internal spreader canals, two internal collection canals, two seepage canals, A2 and Pond 3 Canal; STA perimeter embankment, two interior levees, and a distribution canal levee. The structures associated with the NFSTA include A1 Pump Station, inflow pump station, a seepage canal bridge with box culvert, 2 gated inflow culverts and 4 gated outflow culverts within the STA cells, an emergency by-pass inflow structure with four box culverts, a seepage to inflow canal structure, an emergency by-pass outflow structure with four box culverts, and four gated outflow structures for the NFSTA outflow canal.

Surveys of the North Feeder STA (including associated features) construction footprint should be completed prior to construction to identify invasive and non-native plant and animal species that may be spread by construction activities. These species as well as EDRR species should be treated/removed prior to construction. During the construction phase, surveys and treatment/removal of identified species shall be completed. Once features are constructed, the area should be maintained to prevent establishment of invasive species.

While cattail is considered a native nuisance species within the Everglades, cattails will not be targeted for treatment within the North Feeder STA. Cattails are a desired species within STA's; cattails and other emergent native species such as pickerel weed and bulrush will remain to remove nutrients from the water and improve water quality.

Given the utilization of purple swamphen of existing STAs throughout the Everglades region, early detection and removal efforts should be undertaken to prevent localized establishment of the species in the STA. Early efforts will reduce the proliferation of the species and reduce impact to native wading birds and wetland vegetation. Swamphens are known to show particular fondness for the STAs vegetation and will consume much of the bulrush and other desirable species.

Additionally, newly constructed levees should be routinely surveyed for breeding sites of Burmese pythons, Nile monitors and other invasive aquatic reptiles. Newly created dry features in aquatic systems encourage nesting, providing a pathway for the breeding pockets of invasive reptiles to expand northwest, into areas currently free of these reptiles.

Control of invasive sailfin catfish and blue tilapia prior and during the construction of this project will ensure prime functionality of this STA. Sailfin catfish burrow in embankments and cause cave-ins of levee sides. This damage is evident throughout the region and should be actively managed to prevent similar results. Blue tilapias destroy native aquatic vegetation through consumption and nesting activities. Without prior control, the desirable vegetation in the STA will show difficulty in establishment.

Damaging invasive animals such as feral hogs and green iguanas should be monitored and controlled during the construction phase. These species have been demonstrated to cause infrastructure damage throughout the region. Burrowing, tunneling and rooting behavior have caused structural and substrate issues in water control levees, embankments and canals throughout the region. Initial management and control of these species will result in ensuring continued functionality throughout the life of the project.

North Feeder Canal Plug

Surveys of the project area should be completed prior to construction to identify priority invasive and non-native species that may be spread during construction activities. Such species should be treated prior to the beginning of construction. Once installed the canal plug should be monitored and invasive and non-native species should be treated throughout the construction phase.

The plug location may serve as key choke points that can serve to stymie invasive fish species. Structurally damaging sailfin catfish and blue tilapia should receive focused management at this site during low-water events to prevent further spread throughout the features of this project.

Depending on the methods used for completing the plugs, earth moving species such as feral hogs or iguanas may need to be monitored/controlled to ensure there is no damage to features installed.

F.6.8.2 Region 2

Wingate Mill Canal Plug

Surveys of the project area should be completed prior to construction to identify priority invasive and non-native species that may be spread during construction activities. Such species should be treated prior to the beginning of construction. Once installed the canal plug should be monitored and invasive and non-native species should be treated throughout the construction phase.

The plug location may serve as key choke points that can serve to stymie invasive fish species. Structurally damaging sailfin catfish and blue tilapia should receive focused management at this site during low-water events to prevent further spread throughout the features of this project.

Depending on the methods used for completing the plugs, earth moving species such as feral hogs or iguanas may need to be monitored/controlled to ensure there is no damage to features installed.

Wingate Mill & Lard Can Canal Modifications/Discharge Slough

The Wingate Mill Canal Modifications will include the installation of a weir, canal backfill with embankment degradés (2.4 miles) and a spreader canal with embankment degradés (1,500 feet). The Lard Can Canal Modifications will include canal backfill with embankment degradés (1.5 miles), a

spreader canal (.5 miles), flow diversion berm (1.5 miles) and ditch backfill with embankment degradés (.6 miles). Structures will be installed at the West Boundary Road which will be a gated culvert and four culverts will be installed at Ranch Road.

Surveys of the project area should be completed prior to construction to identify priority invasive and non-native species that may be spread during construction activities. Such species should be treated prior to the beginning of construction. Once installed the features should be monitored and invasive and non-native species should be treated throughout the construction phase.

Focus should be placed on controlling damaging species of wildlife, such as feral hogs, iguana, purple swamphen, sailfin catfish and tilapia. The canals and levee features are at damage risk from these species. Early investment in control during the construction phase can prevent significant utilization by these species, ensuring project functionality.

Further, management and prevention activities targeting Burmese python, Nile monitor, and bullseye snakehead will prevent this feature from being an introduction pathway that would facilitate the spread of these species into unaffected areas.

F.6.8.3 Region 3

L-28i Inline Weir

Surveys of the project area should be conducted prior to installation of the weir. Vegetation that could impact the functionality of the weir should be treated throughout the construction phase and includes floating and submersed species as well as tussocks.

This feature will serve as a focal point for the prevention of invasive fish spread throughout the system. During low water events, targeted management can be conducted to prevent the spread of damaging fish species into features of this project.

L-28i Levee Degrade and Canal Backfill

Surveys of the project area should be completed prior to construction to identify priority species that may be spread by construction activities as well as species that should be treated prior to construction. Such species should be treated prior to the beginning of construction. Periodic surveys of the degraded and backfill areas adjacent to the project site should be conducted throughout the construction phase to identify growth of priority species. Water that is spread as a result of the degrade and backfill may result in the spread of priority species and should thereby be closely monitored. Priority plant species in these areas should be treated. It is recommended that adjacent areas within 0.5 mile of the degrade and backfill be systematically surveyed and treated to eliminate close proximity seed sources, thereby preventing spread of priority plant species, such as Brazilian pepper. Invasive and non-native vegetation present within the vegetation restoration area adjacent to the L-28i extension levee shall be treated/removed utilizing a systematic approach in order to minimize the release of nutrients and impacts south of the area. Once areas have been treated, invasive and non-native vegetation shall be maintained in that area during the entire construction phase.

A focal point of invasive wildlife management during construction should be continual monitoring and removal of nesting Burmese pythons from levees prior to backfill activities. Burmese pythons

have historically concentrated on this levee, seeking high ground to lay eggs during breeding season. As demonstrated by restoration efforts in nearby Picayune Strand, as levees are backfilled, pythons become increasingly concentrated on remaining high ground areas. Once levees are completely degraded, pythons will seek new higher, dryer areas to nest. They will travel incredible distances to accomplish this objective. By removing the pythons during the construction phase, exponentially less pythons will disperse into adjacent federal, state and tribal areas. Python control currently conducted by the Miccosukee Tribe of Indians of Florida (Alligator Alley) Reservation area should be examined for avenues of collaboration to increase management efficiency and success.

Restoration Areas

Surveys of the restoration areas (estimated 4,878 acres and 7,472 acres) should be completed during the beginning of the construction phase to determine the area and vegetation to be treated. It is anticipated that vegetation currently present will expand spatially prior to the beginning of the construction phase and therefore additional acreage of vegetation may need to be treated.

Vegetation within the area identified will be treated incrementally over a five-year period. Initial aerial treatments utilizing herbicides will be completed over 25% of the area identified for treatment each year for four years. The first treatment will be nearest the S-140 or L-28 extension and subsequent treatments proceeding south each year. The approximate annual treatment area will be 1,200 and 1,875 acres for the L-28 Interceptor and S-140 sites, respectively. Follow-up treatments will be conducted following the initial treatment in order to treat vegetation that was not initially treated or new vegetative growth. The follow-up treatments will be ground-based, preferably via airboat in order to minimize ground disturbance. These treatments will begin six months to one year post initial aerial treatment. Various factors such as time of year and water levels will impact the timing of the treatments. It is anticipated that follow-up treatments will continue over a five-year period however vegetation should be maintained throughout Construction and OTMP Phases.

Efforts similar to those mentioned in the L-28i backfills should be conducted during the construction phase of this project. This feature construction will allow for targeted removals of Burmese python, limiting their dispersal to other features within this project, and outside of this. A variety of methods from surveys to radio-telemetry control can be utilized to accomplish this objective. When fully employed, these methods will remove breeding aggregations and prevent python dispersal onto local private, state, federal, and tribal lands.

F.6.8.4 Region 4

L-28(S) Levee Degrade, Canal Backfill and Gated Culverts

Surveys of the project area should be completed prior to construction to identify priority species that may be spread by construction activities as well as species that should be treated prior to construction. Such species should be treated prior to the beginning of construction. Periodic surveys of the degraded and backfill areas adjacent to the project site should be conducted throughout the construction phase to identify growth of priority species. Water that is spread as a result of the degrade and backfill as well as by the new water control structures, may result in the spread of priority species, and should thereby be closely monitored. Priority plant species in these areas should be treated. It is recommended that adjacent areas within 0.5 mile of the degrade and backfill be systematically surveyed and treated to eliminate close proximity seed sources, thereby preventing spread of priority plant species, such as Brazilian pepper.

Efforts similar to those mentioned in the L-28i backfills should be conducted during the construction phase of this project. This feature construction will allow for targeted removals of Burmese python, limiting their dispersal to other features within this project, and outside of this. A variety of methods from surveys to radio-telemetry control can be utilized to accomplish this objective. When fully employed, these methods will remove breeding aggregations and prevent python dispersal onto local private, state, federal, and tribal lands.

Invasive wildlife control that includes focus on burrowing species such as sailfin catfish and iguana is necessary to prevent future damage to the construction site. By addressing these species initially, less effort is needed to control the invasive wildlife in the OMMR&R phase of the project.

L-29 Plug

Surveys of the project area should be completed prior to construction to identify priority invasive and non-native species that may be spread during construction activities. Such species should be treated/removed prior to the beginning of construction.

Depending on the methods used for completing the plug, earth moving species such as feral hogs or iguanas may need to be controlled to ensure there is no damage to features installed. Once installed the canal plug should be monitored and invasive and non-native species should be treated/removed throughout the construction phase.

The plug location may serve as key choke points that can serve to stymie invasive fish species. Structurally damaging sailfin catfish and blue tilapia should receive focused management at this site during low-water events to prevent further spread throughout the features of this project.

Surveys of the L-29 Canal should occur prior to moving water in order to determine if there are invasive species that may be introduced as a result of the connection. Priority invasive species should be treated/removed prior to moving water to the new location. These features should be surveyed during the construction phase to ensure bullseye snakehead, spectacled caiman, and other priority species movements are not facilitated by this project. Surveillance and rapid removal of priority species will help prevent this project from being a source of dispersal on invasive wildlife.

Culverts – 11 Mile Road, US 41 (Tamiami Trail) and Loop Road

Surveys of the project area should be completed prior to construction to identify priority species that may be spread by construction activities. Priority species should be treated/removed prior to construction and culvert installation.

Invasive wildlife control that includes focus on burrowing species such as sailfin catfish and iguana is necessary to prevent future damage to the construction site. By addressing these species initially, less effort is needed to control the invasive wildlife in the OMMR&R phase of the project.

F.6.9 Specific Control by Project Feature – OMRR&R Phase

F.6.9.1 Region 1

North Feeder STA

Surveys of the North Feeder STA (including associated features) construction footprint should be completed on a routine basis to identify invasive and non-native plant and animal species. These species as well as EDRR species should be treated/removed during the OMRR&R Phase. The area should be maintained in order to prevent establishment of invasive species.

While cattail is considered a native nuisance species within the Everglades, cattails will not be targeted for treatment within the North Feeder STA. Cattails are a desired species within STA's; cattails and other emergent native species such as pickerel weed and bulrush will remain to remove nutrients from the water and improve water quality.

Information gathered during construction phase will enable this feature to be placed into an integrated wildlife damage program with area features. The purpose of this would be to control damaging species such as iguana, swamp hen, sailfin catfish and feral hog. Emphasis will be placed on minimizing the impacts that these invasive species will have to the functionality of the STA. Additionally, this program will promote ecological stability by limiting the spread of invasive wildlife into the system.

North Feeder Canal Plug

Surveys of the project area should be completed on a routine basis to identify invasive and non-native plant and animal species. These species as well as EDRR species should be treated/removed during the OMRR&R Phase. The area should be maintained to prevent establishment of invasive species.

During the OMRR&R phase, emphasis should be placed on monitoring for species that may cause damage to the plug. This may come in the form of burrowing fish or feral swine, depending on the feature's construction. Periodic monitoring of the invasive species impact on this specific feature, accompanied by targeted management if needed, is the best procedure for addressing invasive species on this feature.

F.6.9.2 Region 2

Wingate Mill Canal Plug

Surveys of the project area should be completed on a routine basis to identify invasive and non-native plant and animal species. These species as well as EDRR species should be treated/removed during the OMRR&R Phase. The area should be maintained to prevent establishment of invasive species.

During the OMRR&R phase, emphasis should be placed on surveillance for species that may cause damage to the plug. This may come in the form of burrowing fish or feral swine, depending on the feature's construction. Periodic monitoring of the invasive species impact on this specific feature, accompanied by targeted management if needed, is the best procedure for addressing invasive species on this feature.

Wingate Mill & Lard Can Canal Modifications/Discharge Slough

Surveys of the project area should be completed on a routine basis to identify invasive and non-native plant and animal species. These species as well as EDRR species should be treated/removed prior during the OMRR&R Phase. The area should be maintained to prevent establishment of invasive species.

During the OMRR&R phase, damaging wildlife species will be given a priority in management. In order to promote function of the project, invasive fish and wildlife species will be controlled to minimize damage caused to vegetation, embankments and structures. Species such as sailfin catfish and feral hogs will be controlled to ensure structural integrity, while purple swamphen and blue tilapia will be managed to limit beneficial vegetation destruction.

F.6.9.3 Region 3

L-28i Inline Weir

Surveys of the project area should be conducted on a routine basis. Vegetation that could impact the functionality of the weir should be treated throughout the OMRR&R Phase, vegetation to be treated includes floating and submersed species as well as tussocks.

Similar to other features, this weir will best be protected through regular inspection for invasive wildlife damage and utilization. Targeted management efforts can then be undertaken to address any damage caused, species dependent.

L-28i Levee Degrade and Canal Backfill

Periodic surveys of the project area and effected areas adjacent to the project site should be conducted throughout the OMRR&R Phase to identify growth of invasive species. New growth of invasive species in these areas should be treated/removed. The restoration area will require routine maintenance in order to control invasive plant species. Adaptive Management techniques should be implemented to assist with management of invasive plant species in the restoration area.

With the vast majority of invasive reptiles controlled in this feature during the construction phase, surveillance and spot treatments of higher areas such as tree islands can ensure removal of remnant or volunteering individuals.

Restoration Areas

Periodic surveys of the restoration areas should be conducted throughout the OMRR&R Phase to identify growth of invasive species. Any new growth of invasive plant species in these areas should be treated.

Invasive wildlife control centering on structure protection to monitor, manage and reduce risk from sailfin catfish, iguana, and feral hog are necessary for continual operation of these culverts. Periodic management efforts to reduce damaging species in the immediate area of the footprint will prevent cave-ins, obstructions, and maintain waterflow through these culverts.

With the vast majority of invasive reptiles controlled in this feature during the construction phase, surveillance and spot treatments of higher areas such as tree islands can ensure removal of remnant or volunteering individuals.

F.6.9.4 Region 4

L-28(S) Levee Degrade, Canal Backfill and Gated Culverts

Surveys of the project area as well as the restoration areas should be completed on a routine basis to identify invasive and non-native plant and animal species. These species as well as EDRR species should be treated/removed during the OMRR&R Phase. The area should be maintained to prevent establishment of invasive species.

Similar to the L-28i degrade and backfill, surveillance and management of species should be conducted to ensure new habitat does not become reinfested with invasive wildlife. Invasive species that can cause structural damage such as iguanas, feral hogs and sailfin catfish should be controlled to insure proper operation of these features. Periodic surveillance and observation can be followed up with targeted management of problem species if needed.

L-29 Plug

Periodic surveys of the project area and effected areas adjacent to the project site should be conducted throughout the OMRR&R Phase to identify growth of invasive species. New growth of invasive species in these areas should be treated/removed.

Invasive species that can cause structural damage such as iguanas, feral hogs and sailfin catfish should be controlled to insure proper operation of these features. Periodic monitoring and observation can be followed up with targeted management of problem species if needed.

Culverts – 11 Mile Road, US 41 (Tamiami Trail) and Loop Road

Periodic surveys of the project area and effected areas adjacent to the project site should be conducted throughout the OMRR&R Phase to identify growth of invasive species. New growth of invasive species in these areas should be treated/removed.

Invasive wildlife control centering on structure protection to monitor, manage and reduce risk from sailfin catfish, iguana, and feral hog are necessary for continual operation of these culverts. Periodic management efforts to reduce damaging species in the immediate area of the footprint will prevent cave-ins, obstructions, and maintain waterflow through these culverts.

F.7 Education / Outreach Opportunities at Recreational Areas

Recreational opportunities will be created by the LRWRP. Recreation areas such as boat ramps, hiking trails, and hunting areas can serve as vectors and pathways for aquatic and terrestrial invasive species. For example, invasive species can be transferred from one area to another by hikers and by boats/trailers. Many recreational users are unaware of their role in the spread of unwanted species. Hence, educating the public on preventing the spread of invasive species can be a cost-effective component of the overall management strategy. The recreation access points can be used to display educational information on invasive species identification, prevention/control measures, and

awareness of the invasive species programs in the area, and how individuals can contribute to invasive species prevention. Educational kiosks are recommended and should include information on:

- Specific priority invasive species in the area
- Impacts and costs of invasive species on conservation, human health, and recreation
- Preventative measures, such as removing vegetation from boats/trailers before leaving the boat ramp or removing vegetation from shoes and clothing before leaving the area.
- Ways to report invasive species observations
- Programs that citizens can get involved with and learn more about invasive species
- Laws against the release of non-native wildlife

F.8 Costs

A summary of costs are can be found on **Table F-5** with further detailed costs of the construction phase in **Table F-6**, OMRR&R in **Table F-7** and monitoring in **Table F-8**.

It was assumed that in the field baselines and potential invasive species treatments and management would need to occur starting about 2 years prior to the actual construction start date. However, in some cases, it may prove beneficial to begin invasive species management upon the initial start of Construction Phase. Costs were estimated for the life of the project, assuming a 50-year life. However, due to size, the OMRR&R table only shows years 1 and the total 50-year cost estimate. Monitoring costs are provided for pre-construction, OTMP and OMRR&R.

Table F-1. Invasive Plant Species Documented in the Project Area

Common Name	Scientific Name	Region Documented In				FLEPPC Category
		LO	NE	EAA	GE	-
earleaf acacia	<i>Acacia auriculiformis</i>	x	x		x	I
sisal	<i>Agave sisalana</i> Perrine	x	x	x	x	II
woman's tongue tree	<i>Albizia lebbek</i> (L.) Benth	x	x	x	x	I
alligatorweed	<i>Alternanthera philoxeroides</i> (Mart.) Griseb.	x	x	x	x	II
Sprenger's asparagus fern	<i>Asparagus aethiopicus</i> L.	x	x	x	x	
mosquito fern	<i>Azolla pinnata</i>	x			x	
common bamboo	<i>Bambusa vulgaris</i> Schrad.		x		x	
Javanese bishopwood	<i>Bischofia javanica</i> Blume	x	x	x	x	I
paper mulberry	<i>Broussonetia papyrifera</i>	x	x	x	x	II
bottlebrush	<i>Callistemon viminalis</i> (Gaertn.) G. Don ex Loudon	x	x	x	x	II
river sheoak	<i>Casuarina cunninghamiana</i>	x	x	x	x	
Australian-pine	<i>Casuarina equisetifolia</i> L.	x	x	x	x	I
gray sheoak	<i>Casuarina glauca</i> Sieb. ex Spreng	x	x	x	x	I
coconut palm	<i>Cocos nucifera</i>				x	II
coco yam, wild taro	<i>Colocasia esculenta</i> (L.) Schott	x	x	x	x	I
Asian nakedwood	<i>Colubrina asiatica</i> (L.) Brongn.	x	x	x	x	I
smooth crotalaria	<i>Crotalaria pallida</i> Aiton	x	x	x	x	
showy rattlebox	<i>Crotalaria spectabilis</i> Roth	x	x	x	x	
umbrella plant	<i>Cyperus involucratus</i> Rottb	x	x	x	x	II
crowfootgrass	<i>Dactyloctenium aegyptium</i> (L.) Willd	x	x	x	x	II
air-potato	<i>Dioscorea bulbifera</i> L.	x	x	x	x	I
waterhyacinth	<i>Eichhornia crassipes</i> (Mart.) Solms	x	x	x	x	I
Cupid's-shaving-brush	<i>Emilia fosbergii</i> D.H. Nicols.	x	x	x	x	

Common Name	Scientific Name	Region Documented In				FLEPPC Category
		LO	NE	EAA	GE	-
Surinam cherry	<i>Eugenia uniflora</i> L.	x	x	x	x	I
Chinese crown orchid	<i>Eulophia graminea</i>	x	x	x	x	II
Chinese banyan	<i>Ficus microcarpa</i> L. f.	x	x	x	x	I
hydrilla	<i>Hydrilla verticillata</i> (L. f.) Royle	x	x	x	x	I
miramar weed, green swampweed	<i>Hygrophila polysperma</i> (Roxb.) T. Anders.	x	x		x	I
West Indian marsh grass	<i>Hymenachne amplexicaulis</i> (Rudge) Nees	x	x	x	x	I
jaraguagrass	<i>Hyparrhenia rufa</i> (Nees) Stapf	x	x		x	II
cogongrass	<i>Imperata cylindrica</i> (L.) Beauv.	x	x	x	x	I
Lantana, shrub verbena	<i>Lantana camara</i>	x	x	x	x	I
white leadtree	<i>Leucaena leucocephala</i> (Lam.) de Wit	x	x	x	x	II
Asian marshweed	<i>Limnophila sessiliflora</i>	x		x	x	II
primrose-willow	<i>Ludwigia peruviana</i> (L.) Hara	x	x	x	x	I
Old World climbing fern	<i>Lygodium microphyllum</i> (Cav.) R. Br.	x	x	x	x	I
Phasey bean, wild bushbean	<i>Macroptilium lathyroides</i>	x	x	x	x	II
guineagrass	<i>Megathyrsus maximus</i> (Jacq.) R. Webster	x	x	x	x	
melaleuca	<i>Melaleuca quinquenervia</i> (Cav.) Blake	x	x	x	x	I
chinaberry	<i>Melia azedarach</i> L.	x	x	x	x	
natalgrass	<i>Melinis repens</i> (Willd.) Zizka	x	x	x	x	I
plantain	<i>Musa x paradisiaca</i>	x	x	x	x	
Asian swordfern	<i>Nephrolepis brownii</i> (Desv.) Hovenkamp & Miyam.	x	x	x	x	I
narrow swordfern	<i>Nephrolepis cordifolia</i> (L.) C. Presl	x	x	x	x	
burmareed	<i>Neyraudia reynaudiana</i> (Kunth) Keng ex A.S. Hitchc.	x	x	x	x	I
crested floating heart	<i>Nymphoides cristata</i> (Roxb.) O. Ktze.	x	x	x	x	I
monk orchid	<i>Oeceoclades maculata</i> (Cav.)	x	x	x	x	
torpedo grass	<i>Panicum repens</i>				x	I

Common Name	Scientific Name	Region Documented In				FLEPPC Category
		LO	NE	EAA	GE	-
elephant grass, Napier grass	<i>Pennisetum purpureum</i> Schumacher	x	x	x	x	I
Senegal date palm	<i>Phoenix reclinata</i>	x	x	x	x	II
Waterlettuce	<i>Pistia stratiotes</i>	x	x	x	x	I
guava	<i>Psidium guajava</i> L.	x	x	x	x	I
ladder brake	<i>Pteris vittata</i> L.	x	x	x	x	
large flower Mexican clover	<i>Richardia grandiflora</i>	x	x	x	x	II
castorbean	<i>Ricinus communis</i> L.	x	x	x	x	
green shrimp plant, Browne's blechum	<i>Ruellia blechum</i> (var. <i>Blechum brownei</i>)				x	II
water fern	<i>Salvinia minima</i> Baker	x	x	x	x	I
iguanatail	<i>Sansevieria hyacinthoides</i> (L.) Druce	x	x	x	x	II
octopus tree	<i>Schefflera actinophylla</i> (Endl.) H.A.T. Harms	x	x	x	x	I
Brazilian peppertree	<i>Schinus terebinthifolius</i> Raddi	x	x	x	x	I
Brazilian peppertree	<i>Schinus terebinthifolius</i> var. <i>raddianus</i> Engl.	x	x	x	x	I
lakeshore nutrush	<i>Scleria lacustris</i> C. Wright	x	x	x	x	I
Valamuerto	<i>Senna pendula</i>	x	x	x	x	
climbing cassia, Christmas cassia, Christmas senna	<i>Senna pendula</i> var. <i>glabrata</i>	x	x	x	x	I
wetland nightshade	<i>Solanum tampicense</i> Dunal	x	x	x	x	I
tropical soda apple	<i>Solanum viarum</i> Dunal	x	x	x	x	I
annual sowthistle	<i>Sonchus oleraceus</i> L.	x	x	x	x	
Bay Biscayne creeping-oxeye	<i>Sphagneticola trilobata</i> (L.C. Rich.) Pruski	x	x	x	x	
Smutgrass	<i>Sporobolus indicus</i>	x	x	x	x	
cayenne porterweed	<i>Stachytarpheta cayennensis</i> (Rich.) Vahl	x	x	x	x	
St. Augustine grass	<i>Stenotaphrum secundatum</i>	x	x	x	x	
American evergreen, arrowhead vine	<i>Syngonium podophyllum</i> Schott	x	x	x	x	I
Java plum	<i>Syzygium cumini</i> (L.) Skeels	x	x	x	x	I

Common Name	Scientific Name	Region Documented In				FLEPPC Category
		LO	NE	EAA	GE	-
tropical almond	<i>Terminalia catappa L.</i>	x	x	x	x	II
boatlily	<i>Tradescantia spathacea Sw.</i>	x	x	x	x	II
coat buttons	<i>Tridax procumbens L.</i>	x	x	x	x	
Caesarweed	<i>Urena lobata L.</i>	x	x	x	x	I
paragrass	<i>Urochloa mutica (Forsk.) T.Q. Nguyen</i>	x	x	x	x	I
Washington fan palm	<i>Washingtonia robusta</i>		x		x	II

Table F-2. Invasive Plant Species Documented in the Project Area (Totals from Table F-1 above).

	TOTALS
Total Non-native plants	79
Total FLEPPC Category I	39
Total FLEPPC Category II	19
Total Noxious Weeds	13

This list was compiled from the 2021 Florida Exotic Pest Plant Council (FLEPPC) Category I and II species lists. It was cross-checked with species occurrences reported in EDDMapS (Early Detection and Distribution Mapping System) for Broward, Charlotte, Collier, DeSoto, Glades, Hendry, Highlands, Lee, Miami-Dade, Martin, Monroe, Okeechobee, Palm Beach, and St. Lucie counties. Any of the FLEPPC species that had not been recorded in these counties were removed from the list. The list also includes any species that are being actively managed in these areas by the US Army Corps of Engineers or the National Park Service (based on WEEDAR (Weed Data and Reports) data).

Table F-3. Invasive Animal Species Documented in the Project Area.

Common Name	Scientific Name	Region Documented In			
		LO	NE	EAA	GE
BIRDS					
Common Myna	<i>Acridotheres tristis</i>	X	X	X	X
Muscovy Duck	<i>Cairina moschata</i>	X	X	X	X
Rock Dove	<i>Columba livia</i>	X	X	X	X
Hill Myna	<i>Gracula religiosa</i>	X	X	X	X
House Sparrow	<i>Passer domesticus</i>	X	X	X	X
Purple Swamphen	<i>Porphyrio porphyrio</i>	X	X	X	X
Eurasian Collared-Dove	<i>Streptopelia decaocto</i>	X	X	X	X
European Starling	<i>Sturnus vulgaris</i>	X	X	X	X
Sacred Ibis	<i>Threkiornis aethiopicus</i>	X	X	X	X
White-winged Dove	<i>Zenaida asiatica</i>	X	X	X	X
REPTILES & AMPHIBIANS					
Knight Anole	<i>Anolis equestris equestris</i>	X	X	X	X
Cuban Green Anole	<i>Anolis porcatius</i>				X
Brown Anole	<i>Anolis sagrei</i>	X	X		
Brown Basilisk	<i>Basiliscus vittatus</i>	X	X	X	X
Common Boa	<i>Boa constrictor</i>	X	X	X	X
Spectacled Caiman	<i>Caiman crocodilus</i>				X
Veiled Chameleon	<i>Chamaeleo calyptratus</i>	X	X	X	X
Giant Whiptail	<i>Cnemidophorus motaguai</i>				X
Nile Crocodile	<i>Crocodylus niloticus</i>				
Greenhouse Frog	<i>Eleutherodactylus planirostris</i>		X		
Yellow anaconda	<i>Eunectes notaeus</i>			X	X
Common House Gecko	<i>Hemidactylus frenatus</i>				X
Tropical House Gecko	<i>Hemidactylus mabouia</i>	X	X	X	X
Mediterranean Gecko	<i>Hemidactylus turcicus</i>	X	X	X	X
Green Iguana	<i>Iguana iguana</i>	X	X	X	X

Common Name	Scientific Name	Region Documented In			
		LO	NE	EAA	GE
Northern Curlytail Lizard	<i>Leiocephalus carinatus armouri</i>	x	x	x	x
Cuban Treefrog	<i>Osteopilus septentrionalis</i>	x	x	x	x
Giant Day Gecko	<i>Phelsuma madagascariensis grandis</i>				x
Reticulated python	<i>Python reticulatus</i>				x
Burmese Python	<i>Python molurus bivittatus</i>	x	x	x	x
Brahminy Blind Snake	<i>Ramphotyphlops braminus</i>	x	x		x
Giant Toad	<i>Rhinella marina</i>	x	x	x	x
Black and white tegu	<i>Tupinambis merianae Linnaeus, 1758</i>				x
Nile Monitor	<i>Varanus niloticus</i>	x	x	x	x
FISH	-			-	
Oscar	<i>Astronotus ocellatus</i>				x
Pike killifish	<i>Belonesox belizanus</i>	x	x	x	x
Butterfly peacock	<i>Cichla ocellaris</i>				x
Black acara	<i>Cichlasoma bimaculatum</i>				x
Midas cichlid	<i>Cichlasoma citrinellum</i>				x
Jaguar guapote	<i>Cichlasoma managuense</i>	x	x	X	x
Yellowbelly cichlid	<i>Cichlasoma salvini</i>				x
Mayan cichlid	<i>Cichlasoma urophthalmus</i>				x
Walking catfish	<i>Clarias batrachus</i>				x
Grass Carp	<i>Ctenopharyngodon idella</i>	x	x	x	x
Common carp	<i>Cyprinus carpio</i>				x
African jewelfish	<i>Hemichromis letourneuxi</i>	X	x	x	x
Banded cichlid	<i>Heros severus</i>				x
Brown hoplo	<i>Hoplosternum littorale</i>				x
Suckermouth catfish	<i>Hypostomus sp.</i>				
Spotfin spiny eel	<i>Macrogathus siamensis</i>				x
Asian swamp eel	<i>Monopterus albus</i>				x
Blue tilapia	<i>Oreochromis aureus</i>				x
Mozambique tilapia	<i>Oreochromis mossambicus</i>				x

Common Name	Scientific Name	Region Documented In			
		LO	NE	EAA	GE
Nile tilapia	<i>Oreochromis niloticus</i>	X			X
Orinoco sailfin catfish	<i>Pterygoplichthys multiradiatus</i>				X
zebra tilapia	<i>Tilapia buttikoferi</i>				X
Spotted tilapia	<i>Tilapia mariae</i>				X
MAMMALS	-			-	
Coyote	<i>Canis latrans</i>	X	X	X	X
House mouse	<i>Mus musculus</i>				
Norway rat	<i>Rattus norvegicus</i>				
Black rat	<i>Rattus rattus</i>				X
Wild hog, feral pig	<i>Sus scrofa</i>	X	X	X	X
OTHER	-			-	
Asian clam	<i>Corbicula fluminea</i>	X	X	X	X
freshwater jellyfish	<i>Craspedacusta sowerbyi</i>	X	X	X	X
Giant Ramshorn Snail	<i>Marisa cornuarietis</i>	X	X	X	X
Spike-topped applesnail	<i>Pomacea diffusa</i>	X		X	X
Island applesnail	<i>Pomacea insularum (d'Orbigny, 1839)</i>	X	X	X	X
Giant applesnail	<i>Pomacea maculata</i>	X	X	X	X
Fungus (causes laurel wilt)	<i>Raffaelea lauricola</i>				X

Table F-4. Invasive Animal Species Documented in the Project Area (Totals from Table F-3 Above)

-	TOTALS
Total Non-native Animals	69

Table F-5. WERP Invasive and Nuisance Species Management and Monitoring Costs.

	-	Invasive and Nuisance Species Management Costs
2 Year Pre-Construction		\$2,223,623.35
1 Year Pre-Construction		\$1,871,404.38
Construction Phase		\$10,798,987.31
OTMP		\$4,679,873.52
1 Year OMRR&R Phase		\$3,500,329.57
50-Year OMRR&R Phase (Includes Year 1)		\$314,278,175.46
	Total Management Cost	\$333,852,064
	-	Invasive and Nuisance Species Monitoring Costs
2 Year Pre-Construction		\$1,700,001.12
1 Year OTMP		\$1,442,500.00
1 Year OMRR&R Phase		\$1,442,500.00
10 Year OMRR&R Phase		\$14,425,000.00
	Total Monitoring Cost	\$19,010,001.12

Table F-6. Invasive and Nuisance Species Management Costs – Construction PHASE

Feature/Area	Management Activity	Pre-Constructi on 2yrs	Pre-Constructi on 1yr	Construction	OTMP
Region 1					
North Feeder STA (3700 ac)	EDRR/Plant Surveillance	\$ 2,248.16	\$ 2,248.16	\$ 11,240.80	\$ 35,970.56
	Plant Control/Treatment	\$ 403,362.90	\$ 134,454.30	\$ 277,387.70	\$ 667,427.70
	EDRR/ Animal Surveillance	\$ 17,570.00	\$ 17,570.00	\$ 23,426.00	\$ 23,426.00
	Invasive Animal Control	\$ 17,570.00	\$ 17,570.00	\$ 58,565.00	\$ 17,570.00
	Coordination/Inspections/Contract Implementation	\$ 110,187.77	\$ 42,960.62	\$ 92,654.88	\$ 196,347.32
North Feeder Canal Plug (160 a)	EDRR/Plant Surveillance	\$ 1,686.12	\$ 1,686.12	\$ 3,372.24	\$ 3,372.24
	Plant Control/Treatment	\$ 19,380.80	\$ 19,380.80	\$ 3,876.16	\$ 15,504.64
	EDRR/ Animal Surveillance	\$ 5,856.00	\$ 5,856.00	\$ 5,856.00	\$ 5,856.00
	Invasive Animal Control	\$ 5,856.00	\$ 11,713.00	\$ 17,570.00	\$ 17,570.00
	Coordination/Inspections/Contract Implementation	\$ 8,194.73	\$ 9,658.98	\$ 7,668.60	\$ 10,575.72
Region 2					
Wingate Mill Canal Plug (160 ac)	EDRR/Plant Surveillance	\$ 1,686.12	\$ 1,405.10	\$ 4,496.32	\$ 8,992.64
	Plant Control/Treatment		\$ 9,690.40	\$ 11,628.48	\$ 11,628.48
	EDRR/ Animal Surveillance	\$ 5,856.00	\$ 5,856.00	\$ 23,426.00	\$ 11,713.00
	Invasive Animal Control	\$ 17,570.00	\$ 17,570.00	\$ 5,865.00	\$ 17,570.00
	Coordination/Inspections/Contract Implementation	\$ 6,278.03	\$ 8,630.38	\$ 11,353.95	\$ 12,476.03
Wingate Mill/Lard Can Canal Modifications Discharge Slough	EDRR/Plant Surveillance	\$ 5,058.36	\$ 5,058.36	\$ 5,058.36	\$ 20,233.44
	Plant Control/Treatment			\$ 23,620.35	\$ 119,676.44
	EDRR/ Animal Surveillance	\$ 5,856.00	\$ 5,856.00	\$ 17,570.00	\$ 17,570.00
	Invasive Animal Control	\$ 5,856.00	\$ 11,713.00	\$ 17,570.00	\$ 17,570.00
	Coordination/Inspections/Contract Implementation	\$ 4,192.59	\$ 5,656.84	\$ 15,954.68	\$ 43,762.47

Feature/Area	Management Activity	Pre-Constructi on 2yrs	Pre-Constructi on 1yr	Construction	OTMP
Region 3					
L-28i Inline Weir (160 ac)	EDRR/Plant Surveillance	\$ 1,686.12	\$ 1,686.12	\$ 4,496.32	\$ 8,992.64
	Plant Control/Treatment			\$ 3,876.16	\$ 15,504.64
	EDRR/ Animal Surveillance	\$ 5,856.00	\$ 5,856.00	\$ 5,856.00	\$ 5,856.00
	Invasive Animal Control	\$ 5,856.00	\$ 11,713.00	\$ 17,750.00	\$ 11,713.00
	Coordination/Inspections/Contract Implementation	\$ 3,349.53	\$ 4,813.78	\$ 7,994.62	\$ 10,516.57
L-28i Levee Degrade and Canal Backfill (1.5 Mi) - 960 ac	EDRR/Plant Surveillance	\$ 2,248.16	\$ 2,248.16	\$ 2,248.16	\$ 2,248.16
	Plant Control/Treatment		\$ 34,885.44	\$ 34,885.44	\$ 69,770.88
	EDRR/ Animal Surveillance	\$ 52,708.50	\$ 52,708.50	\$ 52,708.50	\$ 11,713.00
	Invasive Animal Control	\$ 245,973.00	\$ 245,973.00	\$ 245,973.00	\$ 46,852.00
	Coordination/Inspections/Contract Implementation	\$ 75,232.42	\$ 83,953.78	\$ 83,953.78	\$ 32,646.01
Region 4					
L-28s Levee Degrade and Backfill (13 mi) - 8320 ac	EDRR/Plant Surveillance	\$ 4,496.32	\$ 4,496.32	\$ 4,496.32	\$ 4,496.32
	Plant Control/Treatment	\$ 201,560.32	\$ 201,560.32	\$ 201,560.32	\$ 201,560.32
	EDRR/ Animal Surveillance	\$ 52,708.50	\$ 52,708.50	\$ 52,708.50	\$ 11,713.00
	Invasive Animal Control	\$ 245,973.00	\$ 245,973.00	\$ 245,973.00	\$ 11,713.00
	Coordination/Inspections/Contract Implementation	\$ 126,184.54	\$ 126,184.54	\$ 126,184.54	\$ 57,370.66
L-29 Pump Station and Discharge Pipes (6.4 mi) 4096 ac	EDRR/Plant Surveillance		\$ 2,248.16	\$ 8,992.64	\$ 4,496.32
	Plant Control/Treatment	\$ 99,229.70	\$ 9,922.97	\$ 327,458.00	\$ 198,459.39
	EDRR/ Animal Surveillance	\$ 5,856.00	\$ 5,856.00	\$ 5,856.00	\$ 5,856.00
	Invasive Animal Control	\$ 17,570.00	\$ 17,570.00	\$ 17,570.00	\$ 17,570.00
	Coordination/Inspections/Contract Implementation	\$ 30,663.92	\$ 8,899.28	\$ 89,969.16	\$ 56,595.43

Feature/Area	Management Activity	Pre-Constructi on 2yrs	Pre-Constructi on 1yr	Construction	OTMP
Culverts- 11mile and Loop Road (6.5 mi) 4,160	EDRR/Plant Surveillance		\$ 2,248.16	\$ 4,496.32	\$ 4,496.32
	Plant Control/Treatment		\$ 10,078.02	\$ 100,780.16	\$ 50,390.08
	EDRR/ Animal Surveillance	\$ 11,713.00	\$ 11,713.00	\$ 11,713.00	\$ 11,713.00
	Invasive Animal Control	\$ 17,570.00	\$ 17,570.00	\$ 29,282.00	\$ 29,282.00
	Coordination/Inspections/Contract Implementation	\$ 7,320.75	\$ 10,402.29	\$ 36,567.87	\$ 23,970.35
Restoration Areas	EDRR/Plant Surveillance	\$ 2,481.60	\$ 22,481.60	\$ 22,481.60	\$ 22,481.60
	Plant Control/Treatment			\$ 6,457,500.00	\$ 1,614,375.00
	EDRR/ Animal Surveillance	\$ 70,000.00	\$ 70,000.00	\$ 70,000.00	\$ 30,000.00
	Invasive Animal Control	\$ 200,000.00	\$ 200,000.00	\$ 200,000.00	\$ 300,000.00
	Coordination/Inspections/Contract Implementation	\$ 73,120.40	\$ 73,120.40	\$ 1,687,495.40	\$ 491,714.15
		Pre-Construction 2 Yrs	Pre-Construction 1 Yr	Construction	OTMP
		\$	\$	\$	\$
		2,223,623.35	1,871,404.38	10,798,987.31	4,679,873.52

Table F-7. Invasive and Nuisance Species Management Costs – OMRR&R Phase

Feature / Area	Management Activity	Year 1 OMRR&R	50-Year OMRR&R
REGION 1			
NORTH FEEDER STA	EDRR/Plant Surveillance	\$ 22,481.60	\$ 3,432,200.31
	Plant Control/Treatment	\$ 338,499.00	\$ 51,677,655.15
	EDRR/ Animal Surveillance	\$ 34,298.01	\$ 5,236,176.64
	Invasive Animal Control	\$ 68,596.02	\$ 5,236,177.16
	Coordination/Inspections/Contract Implementation	\$ 46,387.46	\$ 7,081,838.64
NORTH FEEDER CANAL PLUG	EDRR/Plant Surveillance	\$ 13,488.96	\$ 2,059,320.18
	Plant Control/Treatment	\$ 23,256.96	\$ 3,550,572.26
	EDRR/ Animal Surveillance	\$ 13,719.44	\$ 2,094,506.42
	Invasive Animal Control	\$ 13,719.44	\$ 2,094,506.89
	Coordination/Inspections/Contract Implementation	\$ 6,418.48	\$ 979,890.58
CANALS			
INFLOW/INTAKE/COLLECTION DISTRIBUTION/SPREADER	EDRR/Plant Surveillance	\$ 22,481.60	\$ 3,432,200.31
	Plant Control/Treatment	\$ 77,523.20	\$ 11,835,240.86
	EDRR/ Animal Surveillance	\$ 34,298.01	\$ 5,236,177.16
	Invasive Animal Control	\$ 34,298.01	\$ 5,236,177.16
	Coordination/Inspections/Contract Implementation	\$ 34,298.01	\$ 5,236,177.16
PUMP STATIONS	EDRR/Plant Surveillance	\$ 11,240.80	\$ 1,716,100.15
	Plant Control/Treatment	\$ 9,690.40	\$ 1,479,405.11
	EDRR/ Animal Surveillance	\$ 34,298.01	\$ 5,236,177.16
	Invasive Animal Control	\$ 34,298.01	\$ 5,236,177.16
	Coordination/Inspections/Contract Implementation	\$ 8,952.72	\$ 1,366,785.96
REGION 2			
WINGATE CANAL PLUG	EDRR/Plant Surveillance	\$ 13,488.96	\$ 2,059,320.18
	Plant Control/Treatment	\$ 46,513.92	\$ 7,101,144.52
	EDRR/ Animal Surveillance	\$ 13,719.44	\$ 2,094,506.89

Feature / Area	Management Activity	Year 1 OMRR&R	50-Year OMRR&R
WINGATE MILL/LARD CAN CANAL MODIFICATION DISCHARGE SLOUGH	Invasive Animal Control	\$ 13,719.44	\$ 2,094,506.89
	Coordination/Inspections/Contract Implementation	\$ 34,976.70	\$ 5,339,791.40
	EDRR/Plant Surveillance	\$ 5,058.36	\$ 772,245.07
	Plant Control/Treatment	\$ 54,327.00	\$ 8,293,944.65
	EDRR/ Animal Surveillance	\$ 13,719.44	\$ 2,094,506.89
	Invasive Animal Control	\$ 13,719.44	\$ 2,094,506.89
REGION 3 L-28I INLINE WEIR	Coordination/Inspections/Contract Implementation	\$ 8,369.71	\$ 1,277,778.94
	EDRR/Plant Surveillance	\$ 3,372.24	\$ 514,830.05
	Plant Control/Treatment	\$ 3,876.16	\$ 591,762.04
	EDRR/ Animal Surveillance	\$ 13,719.44	\$ 2,094,506.89
	Invasive Animal Control	\$ 13,719.44	\$ 2,094,506.89
	Coordination/Inspections/Contract Implementation	\$ 346.87	\$ 52,956.06
L-28I LEVEE DEGRADE/CANAL BACKFILL	EDRR/Plant Surveillance	\$ 4,496.32	\$ 686,440.06
	Plant Control/Treatment	\$ 7,752.32	\$ 1,183,524.09
	EDRR/ Animal Surveillance	\$ 34,298.01	\$ 5,236,177.16
	Invasive Animal Control	\$ 6,859.13	\$ 1,047,163.80
	Coordination/Inspections/Contract Implementation	\$ 534.06	\$ 81,533.05
	REGION 4 L-28S LEVEE DEGRADE/BACKFILL	EDRR/Plant Surveillance	\$ 17,985.28
Plant Control/Treatment		\$ 150,000.00	\$ 22,900,062.55
EDRR/ Animal Surveillance		\$ 34,298.01	\$ 5,236,177.16
Invasive Animal Control		\$ 6,859.13	\$ 1,047,163.37
Coordination/Inspections/Contract Implementation		\$ 2,091.42	\$ 319,291.63
L-29 PUMP STATION & DISCHARGE PIPES		EDRR/Plant Surveillance	\$ 8,992.64

Feature / Area	Management Activity	Year 1 OMRR&R	50-Year OMRR&R
	Plant Control/Treatment	\$ 50,000.00	\$ 7,633,354.18
	EDRR/ Animal Surveillance	\$ 6,859.13	\$ 1,047,163.37
	Invasive Animal Control	\$ 34,298.01	\$ 5,236,177.16
	Coordination/Inspections/Contract Implementation	\$ 1,001.50	\$ 152,895.75
CULVERTS- 11 MILE/LOOP ROAD	EDRR/Plant Surveillance	\$ 4,496.32	\$ 686,440.06
	Plant Control/Treatment	\$ 50,000.00	\$ 7,633,354.18
	EDRR/ Animal Surveillance	\$ 34,298.01	\$ 5,236,177.16
	Invasive Animal Control	\$ 13,719.44	\$ 2,094,506.89
	Coordination/Inspections/Contract Implementation	\$ 1,025.14	\$ 156,504.78
RESTORATION AREAS	EDRR/Plant Surveillance	\$ 22,481.60	\$ 3,432,200.31
	Plant Control/Treatment	\$ 1,614,375.00	\$ 26,355,731.90
	EDRR/ Animal Surveillance	\$ 15,000.00	\$ 2,290,006.25
	Invasive Animal Control	\$ 200,000.00	\$ 30,533,416.73
	Coordination/Inspections/Contract Implementation	\$ 18,518.57	\$ 2,827,175.46
	TOTAL 1 YEAR	\$ 3,500,329.57	\$ 314,278,916.19

Table F-8. Invasive and Nuisance Species Management Costs

REGION	MONITORING ACTIVITY	PRE-CONSTRUCTION 2 YEARS	OPERATIONAL TESTING & MONITORING PHASE	OMRR&R - 1 YEAR	OMRR&R - 10 YEARS
REGION 1	DATA MINING	\$58,080	\$0	\$0	\$0
-	FISH & WILDLIFE MONITORING	\$250,000	\$250,000	\$250,000	\$2,500,000
REGION 2	DATA MINING	\$46,464	\$0	\$0	\$0
-	FISH & WILDLIFE MONITORING	\$150,000	\$150,000	\$150,000	\$1,500,000
REGION 3	DATA MINING	\$46,464	\$0	\$0	\$0
-	FISH & WILDLIFE MONITORING	\$150,000	\$150,000	\$150,000	\$1,500,000
REGION 4	DATA MINING	\$69,696	\$0	\$0	\$0
-	FISH & WILDLIFE MONITORING	\$300,000	\$300,000	\$300,000	\$3,000,000
RESTORATION AREAS	DATA MINING	\$61,797	\$0	\$0	\$0
-	FISH & WILDLIFE MONITORING	\$275,000	\$300,000	\$300,000	\$3,000,000
OTHER COST	OVERSIGHT	\$225,000	\$225,000	\$225,000	\$2,250,000
-	COORDINATION	\$11,250	\$11,250	\$11,250	\$112,500
-	ADMINISTRATIVE & CT SUPPORT	\$56,250	\$56,250	\$56,250	\$562,500
-	TOTAL	\$1,700,001	\$1,442,500	\$1,442,500	\$14,425,000

F.9 REFERENCES

- Alexander, G., and J. Marais. 2007. A Guide to the Reptiles of Southern Africa. Cape Town, Struik, 408 p.
- Baber, D. W., and B. E. Coblenz. 1986. Density, home range, habitat use, and reproduction in feral pigs on Santa Catalina Island. *Journal of Mammalogy* 67:512–525.
- Baber, D. W., and B. E. Coblenz. 1987. Diet, nutrition, and conception in feral pigs on Santa Catalina Island. *Journal of Wildlife Management* 51:306–317.
- Barnett D. T., T.J. Stohlgren, C.S. Jarnevich, G.W. Chong, J.A. Ericson, T.R. Davern, and S.E. Simonson. 2007. The art and science of weed mapping. *Environ. Monit. Assess.* 132:235–252.
- Barrett, R. H. 1978. The feral hog at Dye Creek Ranch, California. *Hilgardia* 46:283–355.
- Barrett, R. H., and D. S. Pine. 1980. History and status of wild pigs, *Sus scrofa*, in San Benita County, California. *California Fish and Game.* 67:105–17.
- Belden, R.C. and M.R. Pelton. 1975. European Wild Hog Rooting in the Mountains of Eastern Tennessee. *Proceeding of the Annual Conference of the Southeastern Association of Game and Fish Commissioners*, 29:665-671.
- Bieber, C., and T. Ruf. 2005. Population dynamics in wild boar *Sus scrofa*: ecology, elasticity of growth rate and implications for the management of pulsed resource consumers. *Journal of Applied Ecology* 42:1203–1213.
- Collins, T., J. C. Trexler, L. G. Nico, and T. Rawlings. 2002. Genetic diversity in a morphologically conservative invasive taxon: Multiple swamp eel introductions in the southeastern United States. *Conservation Biology* 16:1024-1035.
- Cowie, R.H. 2013. Biology, systematics, life cycle, and distribution of *Angiostrongylus cantonensis*, the cause of rat lungworm disease. *Hawaii Journal of Medicine and Public Health* 72(2).
- Cuda, J.P. 2009a. Invasive Species: A Florida Perspective. Entomology and Nematology Department, University of Florida, Institute of Food and Agricultural Sciences.
- Cuda, J.P. 2009b. Chapter 8: Introduction to Biological Control of Aquatic Weeds, pp. 55-60. In: *Biology and control of aquatic plants: a best management practices handbook* (Gettys LA, WT Haller and M Bellaud, eds.). Aquatic Ecosystem Restoration Foundation, Marietta GA. 210 pages.
- Dandelot S., C. Robles, N. Pech, R. Verlaque, and A. Caxaubon. 2008. Allelopathic potential of two invasive *Ludwigia* spp. *Aquatic Botany* 88:311-316.
- D’Cruze, N.J. Sabel, K. Green, J. Dawson, C. Gardner, J. Robinson, G. Starkie, M. Vences and F. Glaw. 2007. The first comprehensive survey of amphibians and reptiles at Montagne des Français, Madagascar. *Herpetological Conservation and Biology* 2:87-99.

- Dewey, SA, KA Andersen. 2004. Distinct roles of surveys, inventories, and monitoring in adaptive weed management. *Weed Technology* 18:1449-1452.
- Dorcas, M.E., J.D. Willson, R.N. Reed, R.W. Snow, M.R. Rochford, M.A. Miller, W.E. Meshaka, P.T. Andreadis, F.J. Mazzotti, C.M. Romagosa, K.M. Hart. 2012. Severe mammal declines coincide with proliferation of invasive Burmese pythons in Everglades National Park. *Proceedings of the National Academy of Sciences of the United States of America*; Vol 109 no. 7 pages 2418-2422.
- Doren, R.F., J.C. Volin and J.H. Richards. 2009. Invasive exotic plant indicators for ecosystem 1633 restoration: An example from the Everglades Restoration Program. *Ecological Indicators*, 1634 9S:S29-S36.
- Doren, R.F., A.P. Ferriter, and H. Hastings (Eds.). 2001. *Weeds Won't Wait!: The Strategic Plan for Managing Florida's Invasive Exotic Plants, Part One: An Assessment of Invasive Plants in Florida. A Report to the South Florida Ecosystem Restoration Task Force and Working Group, FL. 273 pp.*
- Dozier, J. 2012 The Mikania micrantha wrap up. Everglades Cooperative Invasive Species Management Area Newsletter 3(1): March.
- Dray, Jr., F.A. and T.D. Center. 1989. Seed production by Pistia stratiotes L. (water lettuce) in the United States. *Aquatic Botany* 33(1-2):155-160.
- Duever, M.J., J.E. Carlson, J.F. Meeder, L.C. Duever, L.H. Gunderson, L.A. Riopelle, T.R. Alexander, R.L. Myers and D.P. Spangler. 1986. *The Big Cypress National Preserve. Research Report 8, National Audubon Society, New York, NY.*
- Ellis, T.M. 1980. *Caiman crocodilus*: An established exotic in South Florida. *Copeia* 1:152-154.
- Elzinga, C.L., D.W. Salzer, J.W. Willoughby. 1998. Measuring and Monitoring Plant Populations. BLM Technical Reference 1730-1.
- Executive Order Number 13112 64 F.R. 6183 (February 8, 1999).
- Ferriter A., Thayer D., Goodyear C., Doren B., Langeland K., and J. Lane. 2005. Chapter 9: Invasive Exotic Species in the South Florida Environment. In: 2005 South Florida Environmental Report. South Florida Water Management District.
- FLEPPC. 2017. Florida Exotic Pest Plant Council's 2017 list of invasive plant species. Florida Exotic Pest Plant Council, <http://www.fleppc.org/>.
- Florida Department of Environmental Protection. Weed Alert: Cogon grass (*Imperata cylindrical*) Florida Department of Environmental Protection, Bureau of Invasive Plant Management, 3900 Commonwealth Blvd, MS 705, Tallahassee, FL 32399.
- Florida Department of Agriculture and Consumer Services. 2011. Laurel Wilt/Redbay Ambrosia Beetle Detection Update. Tallahassee, FL.

- Florida Fish and Wildlife Conservation Commission. (2018) "Green Iguana."
<http://myfwc.com/wildlifehabitats/nonnatives/reptiles/green-iguana/>
- FWC. 2013. Species Profiles: Nonnatives – Veiled Chameleon. Florida Fish and Wildlife Conservation Commission, Tallahassee, FL. Available online at
<http://myfwc.com/wildlifehabitats/nonnatives/reptiles/veiled-chameleon/>.
- Francis, J.K. Brazilian pepper tree. United States Department of Agriculture, Forest Service, International Institute of Tropical Forestry (San Juan, PR) in cooperation with the University of Puerto Rico (Rio Piedras, PR).
- Frank, J.H. and R.D. Cave. 2005. *Metamasius callizona* is Destroying Florida's Native Bromeliads. Pages 91–101 in: M.S. Hoddle (ed.), *Second International Symposium on Biological Control of Arthropods, September 12–16, 2005, Davos, Switzerland, Volume I*. Forest Service Publication FHTET-2005-08, United States Department of Agriculture, Washington, DC.
- Gordon, D.R. and K.P. Thomas. 1997. Florida's invasion by nonindigenous plants: History, screening, and regulation. Pages 21–37 in: D. Simberloff, D.C. Schmitz and T.C. Brown (eds.), *Strangers in Paradise: Impact and Management of Nonindigenous Species in Florida*, Island Press, Washington, DC.
- Haack, R.A. 2003. Intercepted Scolytidae (Coleoptera) at U.S. ports of entry: 1985-2000. *Integrated Pest Management Reviews* 6: 253-282 (2001).
- Haller, W. 2009. Chapter 7: Mechanical Control of Aquatic Weeds, pp. 41-46. In: *Biology and control of aquatic plants: a best management practices handbook* (Gettys LA, WT Haller and M Bellaud, eds.). Aquatic Ecosystem Restoration Foundation, Marietta GA. 210 pages.
- Hanula JL, Mayfield AE, Fraedrich SW, Rabaglia RJ. 2008. Biology and host associations of redbay ambrosia beetle (Coleoptera: *Curculionidae: Scolytinae*), exotic vector of laurel wilt killing redbay trees in the southeastern United States. *Journal of Economic Entomology* 101: 1276-1286.
- Hill, J.E. and C.A. Watson. 2007. Diet of the nonindigenous swamp eel in tropical ornamental aquaculture ponds in West-Central Florida. *North American Journal of Aquaculture* 69:139–146
- Holm, L.G., D.L. Plucknett, J.V. Pancho and J.P. Herberger. 1977. *The World's Worst Weeds: Distribution and Biology*. University Press of Hawaii, Honolulu, HI.
- Holmes, C.W., Robbins, J.A., Reddy, R.A., Neuman, S.A., and Marot, M. The effect of phosphorous-enriched waters on the timing and rate of cattail growth in the northern Florida Everglades: American Geophysical Union (AGU) Spring Meeting, Washington, D.C., May 28-31, 2002.
- Hughes, T. W. 1985. Home range, habitat utilization, and pig survival of feral swine on the Savannah River Plant. Thesis, Clemson University, Clemson, South Carolina, USA.
- UF/IFAS Extension 2017. SS-AGR-342. Original publication date April 2011. Revised November 2011. Reviewed December 2017. Visit the EDIS website at <http://edis.ifas.ufl.edu>.

- Johnson, S. 2007. The Cuban Treefrog (*Osteopilus septentrionalis*) in Florida. Publication WEC 218, Institute of Food and Agricultural Sciences, University of Florida, Gainesville, FL.
- Krysko, K.L., C.R. Gillette, R.M. Reichart, L.P. Nunez, N.T. Coutu, J.A. Wasilewski, K.M. Enge and A.P. Borgia. 2012. Preliminary dietary analysis for the non-indigenous Oustalet's Chameleon, *Furcifer oustaleti* (Mocquard 1894) (Squamata: Chamaeleonidae), in southern Florida. *IRCF Reptiles & Amphibians* 19(4):280-287.
- Kunzer, J.M. and M.J. Bodle. 2008. *Luziola subintegra* (Poaceae: Oryzaceae), New to Florida and the United States. *Journal of the Botanical Research Institute of Texas* 2: 633-636.
- Langeland, K.A., H.M. Cherry, C.M. McCormick, and K.A. Craddock Burks. 2008. 2nd Edition. Identification and Biology of Non-Native Plants in Florida's Natural Areas. IFAS Publication SP 257. University of Florida, Gainesville.
- Langeland, K.A. and K. Craddock Burks. 1998. Identification and Biology of Non-Native Plants in Florida's Natural Areas. IFAS Publication SP 257. University of Florida, Gainesville.
- Langeland K.A. and J. Hutchinson. 2005. Natural Area Weeds: Old World Climbing Fern (*Lygodium microphyllum*). UF/IFAS document SS-AGR-21. Published 2001. Revised 2005. Reviewed 2008.
- Lass, L. W., T. S. Prather, N. F. Glenn, K. T. Weber, J. T. Mundt, and J. Pettingill. 2005. A review of remote sensing of invasive weeds and example of the early detection of spotted knapweed (*Centaurea maculosa*) and babysbreath (*Gypsophila paniculata*) with a hyperspectral sensor. *Weed Sci.* 53:242-251.
- Laycock, G. 1966. *The Alien Animals*. Natural History Press, Garden City, N.Y.
- Maskell, A.J., J.H. Waddle and K.G. Rice. 2003. *Osteopilus septentrionalis*: Diet. *Herpetological Review*, 34:137.
- Mayfield A.E., Thomas MC. 2009. Florida Department of Agriculture and Consumer Services, Division of Forestry, Gainesville, FL; and J.A. Smith, M. Hughes and TJ Dreaden, School of Forest Resource and Conservation, University of Florida, Gainesville. The redbay ambrosia beetle, *Xyleborus glabratus* Eichhoff (Scolytinae: Curculionidae).
- Mazzotti, F.J., T. Center, F.A. Dray, D. Thayer. 2008. Ecological Consequences of Invasion by *Melaleuca quinquenervia* in South Florida Wetlands: Paradise Damaged, not Lost. IFAS Publication SSWEC123. University of Florida, Gainesville.
- Myers, J. H., D. Simberloff, A. M. Kuris, and J. R. Carey. 2000. Eradication revisited: dealing with exotic species. *Trends Ecol. Evol.* 15:316–320.
- National Invasive Species Council. 2008. 2008-2012 National Invasive Species Management Plan. Department of Interior, Office of the Secretary, Washington, DC. 35 pp.
- National Park Service. National Resources Management: Island Apple Snail. US Department of the Interior. Everglades National Park. South Florida Natural Resources Center, 950 Krome Avenue, Homestead, FL 33030.

- Nico, L.G., P. Sharp, and T.M. Collins. 2011. Imported Asian swamp eels (Synbranchidae: Monopterus) in North American live food markets: potential vectors of non-native parasites. *Aquatic Invasions* 6:69-76.
- Onisko, A.L. 2020. Biology and Management of Two Invasive Scleria Species; *Scleria lacustris* and *Scleria microcarpa*. Thesis, University of Florida, Gainesville, FL.
- Perez, E.A., J.A. Coetzee, T. Ruiz Tellez and M.P. Hill. 2011. A first report of water hyacinth (*Eichhornia crassipes*) soil seed banks in South Africa. *South African Journal of Botany* 77(3) 795-800.
- Pimentel, D., R. Zuniga and D. Morrison. 2005. Update on the Environmental and Economic Costs Associated with Alien-invasive Species in the United States. *Ecological Economics*, 52:273–288.
- Pranty, B., K. Schnitzius, H.W. Lovell. 2000. Discovery, origin, and current distribution of the Purple Swampphen (*Porphyrio porphyrio*) in Florida. *Florida Field Naturalist* 28: 1-11.
- Puri, Atul and W.T. Haller. June 2010. Best Management Practices (BMP's) for *Rotala* and *Nymphoides* Control. UF/IFAS Center for Aquatic and Invasive Plants, FWC Task 155-Annual Report.
- Reed, R.N. and G.H. Rodda. 2009. Giant Constrictors: Biological and Management Profiles and an Establishment Risk Assessment for Nine Large Species of Pythons, Anacondas, and the Boa Constrictor. Open-File Report 2009-1202, United States Geological Survey. Washington, D.C.
- Rejmanek, M. and M.J. Pitcairn. 2002. When is eradication of exotic pest plants a realistic goal? Pages 249-253 in C.R. and M.N. Clout, editors. *Turning the tide: the eradication of invasive species*. IUCN SSC Invasive Species Specialist Group, IUCN, Gland, Switzerland and Cambridge, UK.
- Rew, L. J., B. D. Maxwell, and R. Aspinall. 2005. Predicting the occurrence of nonindigenous species using environmental and remotely sensed data. *Weed Sci.* 53:236–241.
- Rice, K.G., J.H. Waddle, M.W. Miller, M.E. Crockett, F.J. Mazzotti, and H.F. Percival. 2011. Recovery of native treefrogs after removal of nonindigenous Cuban treefrogs, *Osteopilus septentrionalis*. *Herpetologica*, 67(2):105-117.
- Roda, A., G. Nachman, S. Weihman, M.Y. Cong and F. Zimmerman. 2016. Reproductive ecology of the giant African snail in south Florida: Implications for eradication programs. *PLoS ONE* 11(11), e0165408. <http://doi.org/10.1371/journal.pone.0165408>
- Rodgers, L., T. Pernas, and S. D. Hill. 2014. "Mapping invasive plant distributions in the Florida Everglades using the digital aerial sketch mapping technique." *Invasive Plant Science and Management* 7: 360–74.
- Schofield, P.J. and L.G. Nico. 2009. Salinity tolerance of non-native Asian swamp eels (Teleostei: Synbranchidae) in Florida, USA: Comparison of three populations and 1798 implications for dispersal. *Environmental Biology of Fishes*, 85:51-59.
- Seward, N.W., K.C. Vercauteren, G. W. Witmer, and R.M. Engeman. 2004. Feral swine impacts on agriculture and the environment. *Sheep and Goat Research Journal* 19:34-40.

- Shafii, B., W. J. Price, T. S. Prather, L. W. Lass, D. C. Thill. 2003. Predicting the likelihood of yellow starthistle (*Centaurea solstitialis*) occurrence using landscape characteristics. *Weed Sci.* 51:748-751.
- Shafland, P.L., K.B. Gestring, and M.S. Sanford. 2010. An assessment of the Asian swamp eel (*Monopterus albus*) in Florida. *Reviews in Fisheries Science* 18(1):25-39.
- Singer, F.J. 2005. Wild pig populations in the national parks. *Environmental Management*, 5:263-270.
- South Florida Environmental Report. 2013. Chapter 7: Status of Nonindigenous Species. South Florida Water Management District.
- South Florida Environmental Report. 2011. Chapter 9: Status of Nonindigenous Species in the South Florida Environment. South Florida Water Management District.
- Springer, M.D. 1977. Ecologic and economic aspects of wild hogs in Texas. Pages 37-46 in G.W. Wood, ed. *Research and management of wild hog populations*. The Belle W. Baruch For. Sci. Inst. of Clemson Univ., Georgetown, S.C.
- Stein, B.A., L.S. Kutner, and J.S. Adams (Eds.). 2000. *Precious Heritage: The Status of Biodiversity in the United States*. Oxford University Press, Oxford, England.
- Sweeney, J. R., J. M. Sweeney, and S. W. Sweeney. 2003. Feral hog. Pages 1164–1179 in G. A. Feldhamer, B. C. Thompson, and J. A. Chapman, editors. *Wild mammals of North America*. Johns Hopkins University Press, Baltimore, Maryland, USA.
- Taylor, Richard B., Eric C. Hellgren, Timothy M. Gabor and Linda M Ilse. 1998. Reproduction of Feral Pigs in Southern Texas. *Journal of Mammalogy* 79(4): 1325-1331.
- Thorbjarnarson, J.B. 1993. Diet of the spectacled caiman (*Caiman crocodilus*) in the Central Venezuelan Llanos. *Herpetologica* 49(1):108-117
- U.S. Army Corps of Engineers and South Florida Water Management District. 2010. *Melaleuca Eradication and Other Exotic Plants Implement Biological Controls, Final Integrated Project Implementation Report and Environmental Assessment*.
- U.S. Congress, Office of Technology Assessment. *Harmful Non-indigenous Species in the United States, OTA-F-565*. Washington, DC: U.S. Government Printing Office, September 1993.
- USDA. 2013. Plant Pest Information. Giant African Snail. United States 2388 Department of Agriculture, Washington, DC. Available online at http://www.aphis.usda.gov/plant_health/plant_pest_info/gas/index.shtml.
- [USDA] USDA Forest Service Remote Sensing Applications Center. 2012. *A weed manager's guide to remote sensing and GIS*. <http://www.fs.fed.us/eng/rsac/invasivespecies/index.htm> Accessed: October 29, 2012.
- United States General Accounting Office. 2000. *Invasive Species. Federal and Selected State Funding to Address Harmful, Nonnative Species, GAO/RCED-00-219*.

- Waddle, J. H., R.M. Dorazio, S.C. Walls, K.G. Rice, J. Beauchamp, 1819 M.J. Schuman, and F.J. Mazzotti. 2010. A new parameterization for estimating co-occurrence of interacting species. *Ecological Applications*, 20(5):1467-1475
- West, B. C., A. L. Cooper, and J. B. Armstrong. 2009. Managing wild pigs: A technical guide. Human-Wildlife Interactions Monograph 1:1–55.
- Wilcove, D.S., D. Rothstein, J. Dubow, A. Phillips, and E. Losos. 1998. Quantifying threats to imperiled species in the United States. *Bioscience*. 48: 607-615.
- Wilde, S.B. 2005. Avian vacuolar myelinopathy linked to exotic aquatic plants and a novel cyanobacterial species. *Environmental Toxicology* 20(3):348-353.
- Wilde, S.B. 2014. *Aetokthonos hydrillicola* gen. et sp. nov.: Epiphytic cyanobacteria on invasive aquatic plants implicated in avian vacuolar myelinopathy. *Phytotaxa* 181(5):243-260.
- Wunderlin, R.P., and B.F. Hansen. 2008. Atlas of Florida Vascular Plants. [S.M. Landry and K.N. Campbell (application development,) Florida Center for Community Design and Research.] Institute for Systematic Botany, University of South Florida, Tampa. (<http://www.plantatlas.usf.edu/>)
- Wunderlin, R. P., B. F. Hansen, A. R. Franck, and F. B. Essig. 2022. Atlas of Florida Plants (<http://florida.plantatlas.usf.edu/>). [S. M. Landry and K. N. Campbell (application development), USF Water Institute.] Institute for Systematic Botany, University of South Florida, Tampa.
- Zhang, L.Y., W.H. Ye, H.L. Cao and H.L. Feng. 2004. *Mikania micrantha* H.B.K. in China—An overview. *Weed Research* 44:42-49.